Study of the Photon Structure via Dijet Production at HERA

by

Dorian Kçira

Study of the Photon Structure via Dijet Production at HERA

Dissertation
zur Erlangung des Doktorgrades
des Fachbereiches Physik
der Universität Hamburg

vorgelegt von

Dorian Kçira

aus Shkodra

Hamburg
2001

Gutachter der Dissertation:	Prof. Dr. R. Klanner
	Prof. Dr. E. Lohrmann
Gutachter der Disputation:	Prof. Dr. E. Lohrmann
	Prof. Dr. B. Naroska
Datum der Disputation:	12.10.2001
Dekan des Fachbereichs Physik und Vorsitzender des Promotionsausschusses:	Prof. Dr. F.-W. Büßer

Abstract

The dependence of the virtual photon (γ^*) structure on the photon virtuality, Q^2, is studied in the kinematic range $0.1 < Q^2 < 10^4$ GeV^2 and $0.2 < y < 0.55$ by measuring the reaction $e^+p \rightarrow e^+ + jet + jet + X$ with the ZEUS detector at HERA using an integrated luminosity of 38.2 pb^{-1}. Events having two jets with transverse energies $E_T^{*\,jet1} > 7.5$ GeV and $E_T^{*\,jet2} > 6.5$ GeV in the final state have been selected using the longitudinally invariant k_T algorithm in the photon-proton hadronic center of mass frame of reference. The dijet cross section has been measured as a function of the fractional momentum of the photon participating in the hard scattering process, x_γ^{OBS}, and Q^2. The ratio of the dijet cross section with $x_\gamma^{OBS} < 0.75$ to that with $x_\gamma^{OBS} > 0.75$ decreases as Q^2 increases. The data are compared with the predictions of LO pQCD using different parton distribution functions of the photon. The measurements can be interpreted in terms of a resolved photon component that falls with increasing Q^2.

Zusammenfassung

Die Abhängigkeit der Struktur des virtuellen Photons (γ^*) von der Photonenvirtualität, Q^2, wurde in dem kinematischen Bereich $0.1 < Q^2 < 10^4$ GeV^2 und $0.2 < y < 0.55$ untersucht. Hierzu wurde die Reaktion $e^+p \rightarrow e^+ + jet + jet + X$ am ZEUS Detektor bei HERA, basierend auf einer integrierten Luminosität von 38.2 pb^{-1}, gemessen. Mit Hilfe des longitudinal invarianten k_T Algorithmusses wurden Ereignisse mit zwei Jets im Endzustand mit transversalen Energien von $E_T^{*\,jet1} > 7.5$ GeV und $E_T^{*\,jet2} > 6.5$ GeV im Referenzsystem des Photon-Proton Schwerpunktsystems ausgewählt. Der Zwei-Jet Wirkungsquerschnitt wurde als Funktion des Impulsanteils des Photons x_γ^{OBS}, das am harten Subprozess teilnimmt, und Q^2 gemessen. Das Verhältniss des Zwei-Jet Wirkungsquerschnittes mit $x_\gamma^{OBS} < 0.75$ zu dem mit $x_\gamma^{OBS} > 0.75$ verringert sich mit zunehmenden Q^2. Die Daten wurden verglichen mit Vorhersagen führender Ordnung der pQCD für verschiedene Partonendichtefunktionen des Photons. Die Messung kann durch ein aufgelöstes Photon interpretiert werden, dessen Anteil sich mit steigendem Q^2 verringert.

Contents

List of Figures

Chapter 1

Introduction

1.1 The Standard Model

According to the Standard Model, the elementary constituents of matter are quarks and leptons. They are fermions, with spin $1/2$.

There are 6 (known) leptons and 6 quarks. The leptons are: electron (e), muon (μ), tau (τ), electron neutrino (ν_e), muon neutrino (ν_μ), tau neutrino (ν_τ). The quarks are: up quark (u), down quark (d), charme (c), strange quark (s), bottom quark (b), top quark (t). For each particle, there exists a corresponding anti-particle with the same mass but opposite sign charge, isospin, baryon and lepton quantum numbers (see table 1.1).

There are four known fundamental interactions/forces in nature. In quantum field theory these forces are mediated by integer spin gauge bosons. The four known interactions are:

1. the Electromagnetic Interaction
   The Electromagnetic Force is the interaction occurring between particles which carry *electrical charge*. The gauge boson that mediates the electromagnetic force is the photon γ. Photons are massless and electrically neutral (they do not carry electrical charge).

2. the Weak Interaction
   The Weak Force acts between fermions (spin $1/2$ particles) and is mediated via the $W^\pm$ and Z^0 gauge bosons. These gauge bosons are massive[1].

3. the Strong Interaction
   The Strong Force couples particles having *strong colour-charge*, or shortly *colour*. There exist three colours. In this case the gauge bosons are the *gluons*. Gluons themselves are massless but carry colour-charges and are subject to the strong interaction.

4. the Gravitational Interaction
   The Gravitational Force is the force which acts between particles with mass

[1] The W-bosons have a mass of 81 GeV and the Z^0 boson 91.2 GeV.

(m). In quantum theories of the gravity, the gravitational field is mediated by a spin two ($s = 2$) boson which is called *Graviton*.

This interaction is the weakest of the four interactions and is long-ranged. At the present energies used in High-Energy Physics, the gravitational force can be neglected.

The combination of the Electroweak Theory with QCD constitutes the *Standard Model*. In the context of this model the 12 elementary fermions (quarks and leptons) are classified in the Standard Model in three families (generations). The masses of the particles change between families but the way the particles interact remains the same (see also table 1.1).

	Families			electric charge	Forces			
					str	em	weak	grav
Quarks	u	c	t	$2/3$	$\times$	$\times$	$\times$	$\times$
	d	s	b	$-1/3$	$\times$	$\times$	$\times$	$\times$
Leptons	ν_e	ν_μ	ν_τ	0	-	-	$\times$	$\times$
	e	μ	τ	-1	-	$\times$	$\times$	$\times$

Masses of Quarks (MeV)					
d	u	s	c	b	t
< 100	< 100	~ 400	~ 1500	~ 5000	~ 174000

Table 1.1 - Generations of fermions in the Standard Model. The electrical charge of the particles is shown and the interaction they undergo. In the second table estimations of the masses of the quarks are given. The $\times$ in the first table mean that the particles are subject to the force and the $-$ means tha the particles are not subject to it.

Quantum Electrodynamics (QED) is the theory of electromagnetic interactions. The results of QED can be calculated using perturbation theory since the electromagnetic coupling constant $\alpha \simeq 1/137$ is much smaller than unity. At larger distances the effective charges are less than the ones at smaller distances. This comes because of screening effects (charges emit and absorb virtual photons which, on their turn, fluctuate in electron-positron pairs - vacuum polarisation). For the present energy scales in HEP experiments α is small enough that the perturbative approach can be applied.

Strong interactions are described by Quantum Chromodynamics (QCD). The mediators of the strong interaction, the gluons, carry colour. Colour plays for the strong interaction the role of the charge in the electromagnetics interactions. In contrast to the electromagnetic interactions, the gluons, the field quanta of the strong interaction, carry colour and, therefore, interact themselves strongly. The (effective) coupling constant for the strong interactions α_s depends on the scale at which the process occurs. Calculation of results using perturbative methods in QCD is not always possible. In fact perturbative QCD (pQCD) cannot be always applied and its use depends also on the concrete experiment (process) for which the calculations

are made. Contrary to the electromagnetic coupling constant α, the strong coupling constant α_s decreases as the scale increases. Some aspects of QCD will be described in this chapter.

The electromagnetic and weak interactions are unified within the electroweak theory by Glashow, Salam and Weinberg [Gl61]. The QCD and the electroweak theory are gauge theories and can be derived from symmetry principles [Lo92]. For the strong interaction, the basic symmetry is $SU(3)$, which is connected to the three degrees of freedom given by the three colour-charges. The electroweak interaction is based on the $SU(2)xU(1)$ symmetry. $SU(2)$ is the non-abelian electroweak-isospin group. $U(1)$ is the Abelian hypercharge group[2].

Experimental evidence by a large number of high energy experiments has shown an overwhelming success of the Standard Model. Fundamental tests of its correctness have been made and precise determinations of its free parameters are done. In spite of these successes, there are open questions concerning the Standard Model.

Some of the problems and still unresolved questions are listed below:

- *Quark Confinement*, the fact that no free quarks are observed has not been demonstrated in a rigorous way from the Standard Model (in QCD).
- QCD does not predict the masses of quarks and leptons.
- The existence of exactly three families of quarks and leptons is not explained from the Standard Model.
- No combination of the General Theory of Relativity and the Quantum Mechanics exists yet. There is a large and unexplained difference between the scales of the Standard Model (~ 100 GeV) and the scale of Gravity, given by the Planck mass of 10^{19} GeV.
- The Standard Model also does not explain the fact that we observe more particles than antiparticles, although they must have been created in equal numbers.
- One of the main unresolved questions in the Standard Model is the existence of the Higgs Bosons. Its existence is related to the generation of elementary particle masses. Gauge Bosons and Fermions obtain their masses by interacting with the vacuum Higgs Field. Associated with this mechanism is the existence of massive scalar particles called Higgs bosons. A clear experimental observation of the Higgs bosons has not yet been made. The LEP experiments at CERN[3] have produced a number of collisions compatible with the production of Higgs particles with a mass of around 115 GeV. However these events were also compatible with other known processes [LE00], so the evidence was not conclusive.

[2]The hypercharge Y is connected to the isospin I_3 and the electrical charge Q by the relation: $Y = 2 \cdot (Q - I_3)$.

[3]CERN - **C**onseil **E**uropéen pour la **R**echerche **N**ucléaire - Switzerland.

- The Standard Model fails to account for the solar and atmospheric neutrino data which strongly indicate the need for neutrino conversions, as would arise from neutrino-mass-induced oscillations [Ne00].

1.2 Deep Inelastic Scattering

Many aspects of the Standard Model have an important experimental input from the Deep Inelastic Scattering (DIS) regime of high energy physics experiments. The first experimental evidence for the structure of the proton came from elastic scattering experiments performed at the High Energy Physics Laboratory at Stanford University in the early 1950s [Ho53].

In the following section the kinematics of ep scattering is described in detail.

1.2.1 Kinematic Variables for Lepton Nucleon Scattering

The scattering of positrons (electrons) off protons is described by the exchange of a gauge boson: γ^* or Z^0 in the case of neutral current interactions and $W^\pm$ in the case of charged current interactions. In this section the kinematic variables will be defined. The ep collision can be seen as a $\gamma^* p$ collision with the lepton emitting the photon. A Feynman diagram for the single photon exchange is depicted in figure 1.1.

The four momenta for the incoming positron and proton are:

$$\begin{aligned} k &= (E_e, 0, 0, -E_e) \\ p &= (E_p, 0, 0, E_p) \end{aligned}$$

where $E_e = 27.5$ GeV and $E_p = 820$ GeV are respectively the energy of the positron and that of the proton beam. The coordinate system is defined such that the incoming protons move in the positive z-direction. Some important (Lorentz invariant) variables for the description of the kinematics of the ep scattering are:

- the center of mass energy squared of the ep system:

$$s = (p + k)^2 \simeq 4E_e E_p \tag{1.1}$$

  where the approximation is made by neglecting the rest masses of the particles.

- the virtuality (negative of the invariant mass squared) of the exchanged boson (photon):

$$Q^2 = -q^2 = -(k - k')^2 \simeq 2E_e E_e'(1 + \cos\theta_{e'}) \tag{1.2}$$

  E_e' and $\theta_{e'}$ are the energy and the angle of the scattered positron. The upper limit of Q^2 is s: $Q^2 \leq s$.



Figure 1.1 - Diagram for single photon exchange in ep scattering. The four momenta are given in brackets.

- The Bjorken variable x is (in the QPM) the fraction of the proton momentum taken by the struck quark:

$$x = \frac{Q^2}{2p \cdot q} \tag{1.3}$$

- the inelasticity y is the relative energy transfer from the positron to the proton in the proton rest frame:

$$y = \frac{p \cdot q}{p \cdot k} \simeq 1 - \frac{E_e'}{E_e}\frac{1 - \cos\theta_{e'}}{2} \tag{1.4}$$

- the center of mass energy squared of the photon-proton ($\gamma^* p$) system

$$W^2_{\gamma^* p} = (q + p)^2 \simeq sy - Q^2 \tag{1.5}$$

- the (absolute) energy transfer from the positron to the proton in the protons rest frame:

$$\nu = \frac{p \cdot q}{m_p} \simeq \frac{Q^2}{2m_p x} \tag{1.6}$$

The variables Q^2, x, and y are not all independent. They are related through the following formula:

$$Q^2 = s \cdot x \cdot y \tag{1.7}$$

1.2.2 Structure Functions

In DIS ep experiments the emitted high Q^2 virtual photons are used to probe the proton structure. The momentum transfer (squared) can be related to the wavelenth of the virtual boson through the Heisenberg uncertainty principle:

$$\lambda \simeq \frac{\hbar}{\sqrt{Q^2}} \tag{1.8}$$

The smaller λ, the higher is the resolution power of the photon. At higher Q^2 the resolution increases and $q\bar{q}$ pairs originating from gluons can be resolved. Processes like the QCD-Compton and Boson-Gluon-Fusion (BGF) become "visible".

The *Deep Inelastic Scattering* regime is the one where Q^2 is much larger than the mass squared of the proton (and $x \neq 1$). In this regime, the proton can be thought of as a group of quasi-free constituents one of which interacts with the photon while the rest (proton remnant) moves unperturbed.

The Born cross section for *Neutral Current* (NC) DIS can be written in terms of the structure functions of the proton F_i as follows [Ha84]:

$$\frac{d^2\,\sigma(e^{\pm}p)}{dx\,dQ^2} = \frac{2\pi\alpha^2}{xQ^4} \cdot \left[\, Y_+ \cdot F_2(x,Q^2) \;-\; y^2 \cdot F_L \;\mp\; Y_- \cdot xF_3(x,Q^2) \,\right] \tag{1.9}$$

where: $Y_{\pm} = 1 \pm (1-y)^2$ and $F_L = F_2 - 2xF_1$. The structure function $F_3(x,Q^2)$ is the parity violating term of the cross section due to the weak interaction. In the case of Z^0 exchange at high Q^2, where Z^0 exchange becomes important ($Q^2 \simeq m^2_{Z^0}$), the cross section has contributions from the photon, the Z^0, and their interference.

The structure functions F_1, F_2, and F_3 can be interpreted in terms of the parton density functions in the proton (hadron).

Within the single-photon approximation, the ep cross section can be written as a product of the photon flux with the (virtual) photon-proton scattering $\sigma^{\gamma^* p}_{tot}$, provided the photon lifetime is longer than its interaction time with the proton. Virtual photons can be transversely or longitudinally polarised. The cross section can therefore be written as a sum of two parts - the longitudinal and the transverse one:

$$\sigma^{\gamma^* p}_{tot} = \sigma_L + \sigma_T \tag{1.10}$$

The structure functions F_i can be related to these transverse and longitudinal cross sections as follows:

$$F_1 = \frac{Q^2}{8x\pi^2\alpha} \cdot \sigma_T\,; \;\; F_2 = \frac{Q^2}{4\pi^2\alpha} \cdot (\sigma_T + \sigma_L)\,; \;\; F_L = \frac{Q^2}{4\pi^2\alpha} \cdot \sigma_L \tag{1.11}$$

The structure functions F_i are defined with respect to the Born cross section in equation 1.9. One has to note that the measured NC cross section changes from that cross section by a factor $1 + \delta_r(y, Q^2)$. These are the QED radiative corrections and have to be taken into account if one wants to extract the structure functions from the measured cross sections.

1.3 Quark Parton Model

Models to describe the proton structure and that of other hadrons, are based on the *Parton Model* of Feynman [Fe69] and the *Quark Model* by Gell-Mann and Zweig [Ge64]. In the parton model the proton is considered to consist of (quasi) free pointlike objects, called partons. Each parton i carries a fraction of the total momentum of the proton: $p_i = \xi_i \cdot p$. p is the total momentum of the proton, p_i is the momentum of parton i, and $0. \leq \xi_i \leq 1$. The inelastic ep cross section is then given as the incoherent sum of (cross sections of) elastic electron parton scatterings.

In the infinite momentum frame of the proton all the transverse momenta are negligible. In this frame the scaling variable x has received a simple interpretation, it can be interpreted as the fractional longitudinal momentum carried by the struck parton, so it is the same as ξ. Indeed, if we denote by ξ the fractional longitudinal momentum carried by the struck parton, neglecting the proton (and parton) masses, we have:

$$0 \simeq m^2 = (\xi p + q)^2 = \xi^2 p^2 + q^2 + 2\xi qp = \xi^2 p^2 - Q^2 + 2\xi qp \Rightarrow$$

$$\xi = \frac{Q^2}{2pq} = x \tag{1.12}$$

Bjoerken and Paschos suggested (1969) that the fermions which made up the proton according to the two models (partons and quarks) were one and the same thing. Thus the model is called the *Quark Parton Model*.

Since in the Quark Parton model the partons are pointlike, an increase in the momentum transfer Q^2 will not make any new details visible. So one would expect the structure functions to be independent of Q^2. This effect is called *scale invariance*, or shortly *scaling*. Scaling was predicted (1968) by Bjorken. In the high energy limit $Q^2 \to \infty$ but $x = Q^2/2pq$ finite, the structure functions depend only on the dimensionless (scaling) variable x:

$$\begin{aligned} F_1(x,Q^2) &\longrightarrow F_1(x) \\ F_2(x,Q^2) &\longrightarrow F_2(x) \end{aligned} \tag{1.13}$$

The scaling behaviour was confirmed by the SLAC measurements[Bl69].

In the infinite momentum frame, due to Lorentz contraction and relativistic time dilatation, the rate at which partons interact with one another is slowed down. During the time the (virtual) photon interacts with the quark, the quark itself is essentially a free particle, not interacting with the other quarks in the proton. In other words, the parton distributions during the ep collision are effectively "frozen", so that only one parton takes part in the interaction. The probability for an additional parton to take part in the interaction is suppressed as $1/(\pi r_p^2 Q^2)$, where r_p is the radius of the proton. Therefore the ep interaction can be seen as an incoherent sum of scatterings of the electron with the quasi-free quarks in the proton.

In the QPM the structure functions F_1 and F_2 are related to the parton distributions functions of the proton. The structure function F_2 is the charge square weighted sum of the first momenta (momenta densities) in x of the parton distribution functions:

$$F_2(x) = \sum_i e_i^2 x f_i(x) \tag{1.14}$$

where the contributions of $q\bar{q}$ pairs from the sea are neglected. The charges e_i are given in units of the proton charge e. The QPM relates also the structure functions F_1 and F_2 to one another:

$$F_1(x) = \frac{1}{2x} F_2(x) \tag{1.15}$$

The above relation is known as the Callan-Gross relation [Ca69]. It implies that the structure function F_L is zero. This can also be proven using general considerations. By applying the helicity conservation in the Breit frame [Ha84] one can prove that the quarks in the QPM can only couple to transversely polarised photons. Therefore, $\sigma_L^{\gamma^* p} = 0$ is zero $\Rightarrow F_L = 0$.

The QPM was very succesful in explaining many results obtained in the ep experiments. This model is though not perfect and some problems became apparent. One of them is the prediction that all the proton momentum is carried by quarks. The experimental data proved this prediction to be wrong. In fact, from measurements:

$$\sum_i \int_0^1 dx\, x f_i(x) \simeq 0.5 \tag{1.16}$$

The surprise was that less than 50% of the proton's momentum is carried by the quarks. That is most of it is carried by neutral particles. Also the fact that no free quarks can be observed experimentally could not be explained by the QPM. In the not so naive parton model Z^0-boson exchange and QCD gluon radiation are considered. In this improved approximation the quark density functions become a function of Q^2 and x, partially losing its intiutive meaning. The structure function F_2 becomes too a function of Q^2 and x and in the sum over quarks the sea quarks and antiquarks are included.

The problem is treated more rigorously in the Quantum Chromodynamics (QCD), the field theory of strong interactions. In the limit $Q^2 \to \infty$ QCD reproduces the QPM.

1.4 QCD

Quantum Chromodynamics (QCD) is the gauge theory of strong interactions of quarks and gluons. QCD is a non-Abelian gauge theory based on the SU(3) colour symmetry group. Colour in QCD is the equivalent of charge in the electromagnetic interactions. The quarks can have one of three colours. They interact by the exchange of gluons, which are electrically neutral but form a colour charge octet. Gluons were experimentally observed through three-jet events at the PETRA collider at DESY in 1979. The gluons themselves interact strongly since they are not colour neutral. A consequence of this property is the *asymptotic freedom*, which states that the interaction strength of two coloured objects decreases as the distance becomes shorter (see 1.17).

Another property of QCD is *confinement*, which keeps quarks bound into colourless hadrons. This prevents the observation of free quarks. The colour degree of freedom and confinement explain why observed hadrons are made either of $q\bar{q}$ or of triple-quark states. These combinations ensure that the hadrons are colourless and have integer electrical charge. In the presence of QCD, the naive QPM picture of hadrons has to be altered to take into account the radiation and absorption of gluons by quarks as well as the radiation of $q\bar{q}$ pairs by gluons.

QCD has properties which make it much more difficult to work with theoretically than electroweak theory. The strong coupling constant α_s is large. This makes the use of perturbation theory very difficult. The effective (strong) coupling constant α_s depends on the scale at which the QCD process occurs, on the number of quark flavours, and on a free scale parameter:

$$\alpha_s(Q^2) = \frac{4\pi}{(11 - 2n_f/3)\ln(\frac{Q^2}{\Lambda^2})} \tag{1.17}$$

Q^2 is the scale at which α_s is probed. Λ is a QCD cut-off parameter. Measurements show that $\Lambda \simeq 200$ MeV (Particle Data Group: [Gr00]). The (known) number of flavours n_f is 6, so $(11 - 2n_f/3)$ is positive. At large scale Q^2, α_s is small and the quarks are quasi-free. In this region of asymptotic freedom perturbative QCD (pQCD) is applicable and predictions on the properties of strong interactions can be made.

1.4.1 Factorization

Hadron-Hadron and Lepton-Hadron interactions are described in QCD as an incoherent sum of the interaction of the constituents/partons (quarks, gluons) from one hadron with those of the other hadron or with the lepton[4]. The distributions of partons bound in hadrons cannot be calculated from first principles.

A separation of the short-ranged (hard processes) and long-ranged (soft-processes) physics is introduced. This separation is called *factorization*. The QCD factorization

[4]The total cross section of the interaction is an incoherent sum of the cross sections of the interactions of the constituents.

theorem [Co85] states that for hard scattering reactions the cross section can be decomposed into the parton densities of incoming particles and the cross section for the hard process (see figure 1.2). In a simplified form, this can be written:

$$\sigma = f_{i/1}(x_1, \mu_F) \otimes \sigma_{i,j}(q, \mu_F, \alpha_s) \otimes f_{j/2}(x_2, \mu_F) \tag{1.18}$$

$f_{i/1}(x_1, \mu_F)$ and $f_{j/2}(x_2, \mu_F)$ are the parton densities of the incoming particles, while $\sigma_{i,j}(q, \mu_F, \alpha_s)$ is the cross section (matrix element) for the hard scattering of the two partons i and j. The convolutions $\otimes$ imply implicit integration over the fractions x_1 and x_2 of the momentum of the colliding particles carried by the partons i and j.

The parton distribution functions are the probability to find a particular quark having a momentum fraction in the range $(x, x + dx)$. So $f_{i/1}(x_1, \mu_F) \cdot dx_1$ is the probability to find the parton i in the first hadron carrying a fraction x_1 of its momentum.

Generally, parton distribution functions give the probability to find partons (quarks and gluons) in a hadron as a function of the fraction x of the hadron's momentum they carry.

In order to use the factorization, a hard scale μ_F - the factorization scale - has to be introduced for separating the short-ranged from the long-ranged physics. The long-ranged (non perturbative) processes are absorbed in the parton distribution functions. The parton distribution functions are assumed to be universal, process independent. Perturbative QCD does not predict them and their values are fixed from experimental results.

On the other hand, the cross section for the hard process is a short-ranged process and can be perturbatively calculated.

Apart from this, pQCD introduces another scale μ_R - the renormalization scale. By introducing the renormalization scale the divergences coming from the calculation of self-energy diagrams (which yield divergent integrals) are absorbed into the definition of long-ranged parton distribution functions (pdfs). Several renormalization schemes are used, the most importants being the *minimal subtraction scheme* - $\overline{\mathrm{MS}}$ and the *deep inelastic scattering scheme* - DIS.

1.4.2 QCD evolution equations

The Q^2 dependence of the parton distribution functions can be calculated within pQCD. The main origin of this dependence is that a quark seen at a certain scale Q_0^2 as carrying a certain fractional momentum of the hadron x_0 can be resolved into more quarks and gluons if we probe it at a higher scale Q^2. The resolved quarks and gluons carry a smaller fractional momentum of the hadron ($x < x_0$). The change of the structure function F_2 with increasing scale is schematically shown in figure 1.3. Thus, if the whole QCD dynamics is included, the structure function F_2 is expected to rise at low x. This is because the low x region is populated by gluons and sea quarks and the quark density is large. The resulting logarithmic dependence of F_2 on Q^2 at fixed x is known as *scaling violation*. At HERA, the structure function F_2 has been measured in a very wide range of Q^2 and x and the scaling violation for low x has been demonstrated. This is shown in figure 1.4.



Figure 1.2 - Factorization: separation of the short-ranged and long-ranged physics. The parton distributions in the hadrons are not perturbatively calculable (long-ranged physics). The hard cross section $\sigma_{ij}(Q^2)$ is perturbatively calculable (short-ranged physics).

The DGLAP equations (DGLAP:Dokshitzer - Gribov - Lipatov - Altarelli - Parisi [Al77]) are a set of $(2n_f + 1)$ coupled integro-differential equations. They can be used to determine the quark and gluon distribution functions for any value of Q^2 if one knows them at one particular value Q_0^2 within the range of applicability of perturbative QCD. These equations are derived by requiring that the structure functions F_1 and F_2 be independent of the choice of the factorization scale μ_F. This condition can be written: $\mu_F^2 (dF_i(x, Q^2)/d\mu_F^2) = 0$, where $i = 1, 2$. The DGLAP equations were derived in the *leading log approximation* - LLA. The terms which give the dominant contributions at large x and Q^2 were summed to all orders. All other terms were neglected.

In a compact form the DGLAP equations can be written as:

$$\frac{\partial}{\partial \ln Q^2} \begin{pmatrix} q \\ g \end{pmatrix} = \frac{\alpha_s(Q^2)}{2\pi} \begin{bmatrix} P_{qq} & P_{qg} \\ P_{gq} & P_{gg} \end{bmatrix} \otimes \begin{pmatrix} q \\ g \end{pmatrix}, \tag{1.19}$$

$q(x, Q^2)$ and $g(x, Q^2)$ are the quark and gluon distributions. The convolution $\otimes$ denotes an integration. The parameters $P_{ij}(x)$ are called splitting functions. They describe the probability to find a parton of type i with given fractional momentum originating from the parton of type j (i, j can be q, g). Given a specific factorization and renormalization scheme, the splitting functions are obtained in pQCD as an expansion series in α_s:

$$\frac{\alpha_s}{2\pi} P_{ij}(x, Q^2) = \frac{\alpha_s}{2\pi} P_{ij}^{(1)}(x) + (\frac{\alpha_s}{2\pi})^2 P_{ij}^{(2)}(x) + ... \tag{1.20}$$



Figure 1.3 - Behaviour of the structure function F_2 with increasing scale. The two top pictures show the case of QPM. The two bottom pictures show the changes when QCD effects are included (from: [Ab98]).

The truncation after the first two terms in the series defines the next to leading order (NLO) DGLAP evolution.

The DGLAP equations are valid as long as the impact of the neglected terms is small. At very low x this is not true any more. At this region another approach is used and the calculations lead to the BFKL (Balitsky - Fadin - Kuraev - Lipatov: [Ku76]) equations. DGLAP deals with the Q^2 evolution and is inadequate at very low x. BFKL deals with the $\frac{1}{x}$ evolution and is inadequate at large Q^2.

The inclusive measurement of F_2 at HERA has shown that the evolution of structure functions through the DGLAP equations is in good agreement with the experimental results (figure 1.4). Until now it has not been possible to discriminate between



Figure 1.4 - Scaling violations - the structure function F_2 versus the scale Q^2 for different values of x. The structure function F_2 is not independent of Q^2. The F_2 measurements for different x have been scaled by constant factors in order for all of them to fit on the same plot.

DGLAP and BFKL equations. The differences could show up in the exclusive measurements such as high p_T forward jets and in the forward energy flow.

Attempts have been made to achieve a unified DGLAP / BFKL description. Evolution equations which allow an evolution in Q^2 as well as in x are considered in the CCFM evolution equations.

1.5 Photoproduction

In contrast with the high Q^2 regime which has been defined as DIS, the kinematical regime where Q^2 is very small ($Q^2 \simeq 0$) is called *Photoproduction*. In the photoproduction events the scattered electron has a very small angle with respect to the incoming one. Therefore, most of the time it goes undetected in the beam pipe[5]. The exchanged photons in photoproduction are quasi real.

In photoproduction the hard scale of the interactions is not given by Q^2 since Q^2 is very small. In this case the hard scale is given by the transverse energy of the (hard) outgoing partons. The situation now is reversed and the proton is not anymore being probed by the photon. In fact, the photon is being probed by the hard partons from the proton.

This will be discused in more detail in the section 2.4.1.

[5]At HERA, in some cases the scattered electron can be detected in the luminosity detector (Lumi-Electron)

Chapter 2

Introduction to Photon Structure

2.1 Photon Structure

In the context of the Standard Model, the photon is a fundamental, spin one, gauge boson which mediates the electromagnetic interaction. Photons have no mass, no intrinsic structure and no self couplings. Therefore, the concept of the photon structure seems at first paradoxical.

It is well known that the photons couple to particles that carry charge. Hence, photons can fluctuate to fermion-antifermion pairs. These fluctuations, at high photon energies, can live long enough so that the photon interactions with matter can proceed via the interactions of the fermion-antifermion pairs with matter. In such cases the photon interactions can be conveniently described using the concept of photon structure which is similar to (and borrowed from) the concept of the proton structure.

Fundamental differences between photon and proton structures arise from the fact that the photon structure is completely driven from the quantum fluctuations of the photon to fermion-antifermion pairs whereas a considerable part of the proton structure arises from the contributions of the valence quarks which are absent in the case of the photon structure. Therefore, the photon structure functions, unlike those of the proton, exhibit scaling violations already at the quark-parton-model (QPM) level without any need for gluon emission or absorption diagrams.

Measurements of the photon structure allow tests of QED and QCD (depending if the photon fluctuates to charged leptons or quarks). One can therefore speak of “electromagnetic” and “hadronic” structure of the photon.

2.1.1 Hadronic Photon Structure

Photon fluctuations to $q\bar{q}$ pairs are described by QCD [Sc93]. According to QCD, the final state $q\bar{q}$ can radiate gluons, which can further produce quarks. The collection of all final state quarks and gluons, the “quark-gluon cloud”, represent the photon structure.

In this picture, the physical photon can be seen as a superposition of a bare (pointlike) photon (γ_B), refering to the case where the photon does not fluctuate but remains a pointlike photon, and a resolved component (R), where the photon fluctuates:

$$| \gamma > \; = \; Z_3^{1/2} \, | \gamma_B > \; + c \cdot \alpha^{1/2} \, | R > \tag{2.1}$$

Z_3 and $c \cdot \alpha$ are normalization constants. The state $| R >$ must have the same quantum numbers as the photon.



Figure 2.1 - A photon can be bare (pointlike) or fluctuate into a $q\bar{q}$ pair. Low virtuality fluctuations are give rise to vector meson states. From top to bottom, the three components of the photon: bare, anomalous, and VMD component.

The lifetime of the fluctuation $\gamma \to q\bar{q}$ pair is given by the Heisenberg uncertainty principle as:

$$\Delta t \leq \frac{\hbar}{\Delta E} \tag{2.2}$$

where $\Delta E = E_\gamma - E_{q\bar{q}}$ is the energy difference of the photon and the quark-antiquark pair. The virtuality of the $\gamma \to q\bar{q}$ is characterized by the common p_T^2 of the quark and antiquark with respect to the photon direction. Depending on the lifetime of the fluctuations of the photon in $q\bar{q}$, one can distinguish two components of $| R >$. When the virtuality of the photon is small ($\Rightarrow \Delta E$ small) the fluctuation is long-lived. There is then time for a cloud of soft gluons to develop around the $q\bar{q}$ pair, which

forms in this case a bound state with the same quantum numbers as the photon. This process is non-perturbative and the description of this component is based on phenomenology. The vector meson dominance model (VMD) [Sa69] is usually used to describe the non-perturbative component of the hadronic fluctuations of the photon.

In the vector meson dominance model the photon fluctuates into a vector meson: $\gamma \leftrightarrow V$. The lifetime of such a fluctuation is given by:

$$\tau_\gamma^V \simeq \frac{2\hbar \cdot E_\gamma}{m_V^2 + Q^2} \tag{2.3}$$

where E_γ is the energy of the photon, m_V is the mass of the vector meson, and Q^2 is the photon virtuality [1]. The lifetime of the vector meson state becomes shorter with increasing photon virtuality and dies out as $\frac{1}{m_V^2+Q^2}$ with Q^2.

Since the vector meson must have the same quantum numbers as the photon, it has to be a flavour-diagonal vector meson[2]. In the simplest formulation, the VMD asserts that the sole hadronic components of the photon are the three light vector mesons ρ^0, ω, and ϕ. For the VMD component of the photon we could then write:

$$| \gamma_{VDM} > = \sum_{\rho^0,\, \omega,\, \phi} \frac{e}{f_V} \, | V > \tag{2.4}$$

Here $(\frac{e}{f_V})^2$ are the probabilities for the transition of the photon into the respective vector meson. These coefficients are based on low-energy fits and are assumed to be energy independent. Numerically: $f_V^2/4\pi = 2.20,\ 2.36,\ 18.4$ respectively for $\rho^0,\ \omega,\ \phi$. In the generalized vector meson dominance model (GVD) also vector mesons with higher masses are included in the above sum.

When the virtuality is large ($\Rightarrow \Delta E$ large), the fluctuation is short-lived and no bound hadronic state can develop. In this case, a perturbative description is possible. This is the so called anomalous photon component ("pointlike" photon structure), which originates from the perturbative splitting $\gamma \to q\bar{q}$ and the subsequent QCD evolution. This component can be written as:

$$| \gamma_{perturbative} > = \frac{e}{f_{q\bar{q}}} \, | q\bar{q} > \tag{2.5}$$

[1]Lets denote by $q = (E_\gamma, \vec{q})$ the four-momentum of the photon and by $V = (E_V, \vec{p}_V)$ that of the vector meson. Because of momentum conservation $\vec{q} = \vec{p}_V$. The virtuality is $Q^2 = -q^2 = \vec{q}^2 - E_\gamma^2 = \vec{p}_V^2 - E_\gamma^2$. Therefore $\vec{p}_V^2 = Q^2 + E_\gamma^2$.

From the Heisenberg Principle: $\Delta E \cdot \tau_\gamma^V \simeq \hbar$, where τ_γ^V is the lifetime of the fluctuation of the photon in a vector meson and ΔE is energy difference between the two states: $\Delta E = E_V - E_\gamma = \sqrt{\vec{p}_V^2 + m_V^2} - E_\gamma$ $\Rightarrow \Delta E = \sqrt{E_\gamma + Q^2 + m_V^2} - E_\gamma$. The vector meson model is valid for photons of small virtualities (Q^2 small). On the other hand, the mass of the vector meson is much smaller than its energy. So $E_\gamma \gg Q^2 + m_V^2$. With this assumption: $\Delta E \simeq \frac{Q^2+m_V^2}{2E_\gamma} \Rightarrow \tau_\gamma^V \simeq \frac{2\hbar \cdot E_\gamma}{m_V^2+Q^2}$.

[2]As the spin has to be unity and the charge 0

Here $(\frac{e}{f_{q\bar{q}}})^2$ is the probability of the transition of the photon in the quark-antiquark pair.

In total, the photon has three possible states: bare (pointlike), Vector-Meson, and perturbative (anomalous). In general one can finally write:

$$|\gamma> \; = \; Z_3^{1/2}\,|\gamma_B> \; + \frac{e}{f_{q\bar{q}}}\,|q\bar{q}> \; + \sum_{\rho^0,\,\omega,\,\phi} \frac{e}{f_V}\,|V> \tag{2.6}$$

where, for a proper normalization, $Z_3 + (\frac{e}{f_{q\bar{q}}})^2 + \sum_{\rho^0,\,\omega,\,\phi} (\frac{e}{f_V})^2 = 1$. These three components of the photon are represented schematically in figure 2.1.

Most of the time, the photon is a pointlike particle ($Z_3 \simeq 1$). But since the hadronic cross section are very large the (γ^* - hadron) cross sections are dominated by the VMD and anomalous components.

2.2 Photon Structure in e^+e^- experiments - F_2^γ

Originally the photon structure has been studied in e^+e^- experiments. In e^+e^- scattering two photons are emitted from the positron and the electron which interact with one another so that the process can be seen as $\gamma\gamma^{(*)}$ scattering. The process is therefore sensitive to the structure of the photon. The first measurements of the photon structure function were made at PETRA / DESY in Hamburg [PL84]. One of these measurements from the PLUTO collaboration is shown in figure 2.2.

The underlying kinematics in e^+e^- interactions is shown in figure 2.3. The initial state consists of the incoming electron and positron. Each of them emitts a (virtual) photon. The following invariants can be constructed:

- the virtualities of the photons calculated as (minus) squared momenta of them:

$$\begin{aligned} Q^2 &= -q^2 = -(k-k')^2 &= 4E_0E_1'\sin^2(\theta_1'/2) \\ P^2 &= -p^2 = -(l-l')^2 &= 4E_0E_2'\sin^2(\theta_2'/2) \end{aligned} \tag{2.7}$$

  What is called P^2 and Q^2 is a matter convention. Usually Q^2 is the highest virtuality and P^2 the smallest. The photon with the highest virtuality is called the "probe photon" while the photon with the smallest virtuality is called the "target photon".

- The center of mass energy squared of the $\gamma\gamma$ system:

$$W_{\gamma\gamma}^2 = (p+q)^2 \tag{2.8}$$

- The Bjorken variable is denoted here with x_γ. It is the fractional momentum of the photon taking part in the hard interaction:

$$x_\gamma = \frac{Q^2}{2p\cdot q} = \frac{Q^2}{Q^2 + W_{\gamma\gamma}^2 + P^2} \tag{2.9}$$



Figure 2.2 - Measurement of photon structure in e^+e^- experiments. The $F_{2,\mathrm{eff}}^\gamma$ versus x.

- $$y = \frac{p\cdot q}{p\cdot k} \tag{2.10}$$

Three event classes can be distinguished depending on whether the final state electron is tagged or not.

1. If both electrons are not tagged - *anti-tagged / untagged* events - both virtualities Q^2 and P^2 are small and the two photons are considered to be quasi-real. The smallness of the virtualities depends on the detector acceptance.

2. One of the electrons is tagged - *single-tagged* events. In this case, the process of figure 2.3 can be considered as scattering of an electron off a quasi real photon. The quasi real photon has a virtuality $P^2 \simeq 0$ and the hard scale is the virtuality of the tagged photon $Q^2 > 0$. In this case, the structure of the quasi real photon can be probed from the virtual photon in a similar way to the ep scattering (HERA) where the proton structure is probed using a virtual photon.

3. Both electrons are tagged - *double-tagged* events. In this case both virtualities can be measured and a study of the photon structure for different virtualities P^2 is possible. Given the difficulty of doing double-tagged measurements, the statistics for this class of events is poor in comparison to the other two classes above.

The cross section for the reaction $ee \longrightarrow eeX$ can be written:



Figure 2.3 - Kinematics of two-photon induced processes in e^+e^- collisions. k/k' and l/l' are the four-momenta of the incoming/outgoing electrons (positrons). The four momenta are written in brackets for all the components. The photon with the smallest virtuality (P^2) is the target photon while the one with the highest virtuality Q^2 is the "probe photon".

$$\sigma_{ee\to eeX} = \sigma_{e\gamma\to eX} \cdot f_\gamma^e \tag{2.11}$$

The flux factor f_γ^e denotes the flux of the target photon from the incoming electron. This flux takes into account the momentum spread of the target (quasi-real) photon. The calculation of this flux is carried out using the Equivalent Photon Approximation - EPA (see also section 2.4.3).

Since the target photon is almost real ($P^2 \simeq 0$), in most cases it has (almost) only transverse polarization. The $e\gamma \longrightarrow eX$ cross section has only two components:

$$\frac{d^2\sigma(e\gamma \to eX)}{dx_\gamma\, dQ^2} = \sigma_{TT} + \sigma_{LT} \tag{2.12}$$

The first index of the cross sections on the right side of the above relation refers to the polarization of the probing photon. This can be either transverse or longitudinal (T or L) since this photon is virtual. The second index refers to the polarization of the

target (quasi-real) photon and can be only transverse (T). Similarly to the relations (1.11), one can define the photon structure functions using the definitions:

$$\begin{aligned} F_1^\gamma(x_\gamma, Q^2) &= \frac{Q^2}{4\pi^2\alpha}\frac{1}{2x_\gamma}\cdot\sigma_{TT} \\ F_2^\gamma(x_\gamma, Q^2) &= \frac{Q^2}{4\pi^2\alpha}\cdot(\sigma_{TT}+\sigma_{LT}) \\ F_L^\gamma(x_\gamma, Q^2) &= F_2^\gamma(x_\gamma, Q^2) - 2x_\gamma F_1^\gamma(x_\gamma, Q^2) \end{aligned} \tag{2.13}$$

The function F_L^γ is the longitudinal structure function. With these definitions, the $e\gamma$ cross section can be rewritten:

$$\frac{d^2\sigma(e\gamma \to eX)}{dx_\gamma\, dQ^2} = \frac{2\pi\alpha^2}{x_\gamma Q^4} \cdot \left[\left(1 + (1-y)^2\right) \cdot F_2^\gamma(x_\gamma, Q^2) \; - \; y^2 \cdot F_L^\gamma(x_\gamma, Q^2) \right] \tag{2.14}$$

Note the similarity with the relation (1.9) for the ep cross section. Here there is no F_3-part since we only have a photon exchange and have ignored the Z^0 exchange.

The inelasticity y can be reconstructed from the tagged electron as follows: $y = 1 - (E_{tag}/E_0) \cdot \cos^2(\theta_{tag}/2)$, where E_0 is the energy of the incoming electron (the beam energy). E_{tag} and θ_{tag} correspond to the above E_1' and θ_1'. Usually the energy of the tagged electron is large enough so that y is much less than unity (a cut $E_{tag} > E_0/2$ is often applied to reject beam-gas events). Therefore, the y^2F_L term of the $e\gamma$ cross-section can be neglected. In this approximation, the $\frac{d^2\sigma(e\gamma\to eX)}{dx_\gamma\, dQ^2}$ is directly proportional to $F_2^\gamma(x_\gamma, Q^2)$.

In the case of double-tagged events, the target photon is not quasi-real and can have also transverse polarization. In this case additional terms should be considered in the expression (2.12) for the $e\gamma$ cross-section (σ_{TL}, σ_{LL}, cross terms). The simplifications made above do not hold anymore for this class of events and the F_2^γ contribution to the cross section changes. In measuring $F_2^\gamma(x_\gamma, Q^2)$ one has to correct for the fact that P^2 is not zero.

In leading order (LO) the photon structure function F_2^γ can be expressed as a sum over the charge weighted quark densities of the photon:

$$F_2^\gamma(x_\gamma, Q^2) = \sum_i e_i^2 x_\gamma f_i^\gamma(x_\gamma) \tag{2.15}$$

This concept of the structure function F_2^γ has been developed in analogy with the structure function F_2 of the proton (nucleon), although there are some differences between the two:

* The photon has no valence quarks.
* $F_2^\gamma(x_\gamma, Q^2)$ increases with Q^2 for all x_γ. This scaling violation is expected already within the parton model and it comes from the pointlike anomalous contribution even without introducing gluon radiation.



Figure 2.4 - F_2^γ measurements at e^+e^- experiments in different bins of x_γ and Q^2. Note: the variable x on the plots is denoted in the text of the thesis by x_γ. The Q^2 increases from left to right and from top to bottom. The vertical error bars represent the statistical + systematic errors added in quadrature. The horizontal error bars (!) represent the statistical errors only. The mean Q^2 in GeV^2 of each bin is shown also after the experiment mean.

* $F_2^\gamma(x_\gamma, Q^2)$ increases towards $x_\gamma = 1$ while $F_2^{proton}(x, Q^2)$ decreases at large x.

Measurements of F_2^γ versus x_γ (in the plots, this is denoted simply by x) for Q^2 bins in the range $0.2-400$ GeV^2 from different experiments are shown in figure 2.4. If the photon were purely hadronic-like, a steep rise of F_2^γ at low x_γ would be observed, similarly to what HERA has observed for the proton-F_2 at low x. The systematics dominated errors of these measurements are too large to draw exact conclusions. The main systematic problem in extracting F_2^γ is the final-state modeling. A rise at

small x_γ cannot be excluded.

The photon structure function can be written as a sum of two parts:

$$F_2^\gamma = F_2^{\gamma,pointlike} + F_2^{\gamma,hadronic} \tag{2.16}$$

$F_2^{\gamma,hadronic}$ takes into account the vector meson and anomalous contributions. At large x_γ and asymptotically large Q^2, the $F_2^{\gamma,hadronic}$ can be calculated perturbatively (the vector meson part dies out, the anomalous - perturbative contribution remains). In this region one can parametrize this function as [Wi77]:



Figure 2.5 - F_2^γ evolution with Q^2. The $\ln(Q^2)$ dependence over a large range of Q^2 values can be seen. Note: the variable x on the plots is denoted in the text of the thesis by x_γ.

$$F_2^\gamma \sim \frac{a(x_\gamma)}{\alpha_s(Q^2)} + b(x_\gamma) \sim a_1(x_\gamma) \cdot \ln(\frac{Q^2}{\Lambda^2}) + b_1(x_\gamma) \tag{2.17}$$

See also the relation (1.17). a, b, a_1, b_1 are functions of x_γ independent of Q^2. The above relation could be used to extract the QCD scale Λ. At $P^2 \simeq 0$ where the contribution from the non perturbative VDM component is expected to be significant, this is no longer possible.

The Q^2 dependence of F_2^γ is shown in figure 2.5. As can be seen, the results are in agreement with the expected $\ln(Q^2)$ behaviour over two orders of magnitude in Q^2. While the Q^2 dependence is predicted in QCD, the x-dependence is much less well constrained.

The class of anti-tagged events is somehow similar to photoproduction at HERA. Jet production (jet cross sections, jet shapes, etc.) is studied in this regime.

2.3 Parton Distribution Functions of the Photon

Similarly to the case of the proton, parton distribution functions (pdfs) are defined also for the photon. A factorization scale μ_F is introduced and all the long range QCD effects which take part below this scale are absorbed in the parton distribution functions of the photon. The short range processes are calculable in QCD (pQCD).

The parton distribution functions are considered to be universal and are determined from the experiment. The Q^2 dependence of the evolution of the parton distribution functions is however predicted by pQCD.

2.3.1 The evolution of the pdfs of the photon

The evolution equations for the parton densities can be written [Vo97]:

$$\begin{aligned}
\frac{dq_i^\gamma}{d\ln Q^2} &= \frac{\alpha}{2\pi}\overline{P}_{q_i\gamma}\otimes\Gamma^\gamma + \frac{\alpha_s}{2\pi}\Big\{2\sum_{k=1}^{n_f}\overline{P}_{q_iq_k}\otimes q_k^\gamma + \overline{P}_{q_ig}\otimes g^\gamma\Big\}\\
\frac{dg^\gamma}{d\ln Q^2} &= \frac{\alpha}{2\pi}\overline{P}_{g\gamma}\otimes\Gamma^\gamma + \frac{\alpha_s}{2\pi}\Big\{2\sum_{k=1}^{n_f}\overline{P}_{gq_k}\otimes q_k^\gamma + \overline{P}_{gg}\otimes g^\gamma\Big\}\\
\frac{d\Gamma^\gamma}{d\ln Q^2} &= \frac{\alpha}{2\pi}\overline{P}_{\gamma\gamma}\otimes\Gamma^\gamma + \frac{\alpha}{2\pi}\Big\{2\sum_{k=1}^{n_f}\overline{P}_{\gamma q_k}\otimes q_k^\gamma + \overline{P}_{\gamma g}\otimes g^\gamma\Big\}\,.
\end{aligned} \tag{2.18}$$

where $q^\gamma(x_\gamma, Q^2)$ and $g^\gamma(x_\gamma, Q^2)$ are the quark and gluon distributions in the photon and n_f is the number of active flavours. $\overline{P_{ij}}$ are some generalized splitting functions. The difference between the above equations and the DGLAP equations (1.19) is the presence of the inhomogenous terms connected to the pointlike component of the photon $\Gamma^\gamma(x, Q^2)$. This part does not exist in the case of the proton.

Due to charge conjugation invariance and charge conservation, the antiquark distributions are the same as the quark ones: $\bar{q}_i^\gamma(x_\gamma, Q^2) = q_i^\gamma(x_\gamma, Q^2)$ The singlet quark distributions are then $\Sigma^\gamma = 2\cdot\sum_i^{n_f} q_i^\gamma$. The parton densities are subject to the energy-momentum sum rule:

$$\int_0^1 dx_\gamma\, x_\gamma\big[\Sigma^\gamma(x_\gamma, Q^2) + g^\gamma(x_\gamma, Q^2) + \Gamma^\gamma(x, Q^2)\big] = 1 \tag{2.19}$$

If the calculations are restricted to the first order of the QED coupling constant α ($\alpha \ll 1$), the generalized splitting functions reduce to the usual QCD ones $P_{ij}(x, \alpha_s)$. In this case $P_{\gamma q_i}$ and $P_{\gamma g}$ drop out completely. The third row of the equations (2.18) is in this case very simple and can be immediately integrated. Since the $P_{\gamma\gamma} \sim \delta(1-x_\gamma)$ (the density of the photon in the photon), then the LO expression for Γ^γ will be

$$\Gamma_{LO}^\gamma(x_\gamma, Q^2) = \delta(1-x_\gamma)\Big[1 - \frac{\alpha}{\pi}\Big(\sum_q e_q^2 \ln\frac{Q^2}{Q_0^2} + c_1\Big)\Big] \tag{2.20}$$

c_1 is a constant. The momentum sum rule holds order by order in α. Using the above expression, the momentum sum rule becomes:

$$\int_0^1 dx_\gamma\, x_\gamma\big[\Sigma_{LO}^\gamma(x_\gamma, Q^2) + g_{LO}^\gamma(x_\gamma, Q^2)\big] = \frac{\alpha}{\pi}\Big(\sum_q e_q^2 \ln\frac{Q^2}{Q_0^2} + c_1\Big) \tag{2.21}$$

The integrated quark and gluon densities do not sum to unity. Instead they logarithmically increase with Q^2. This comes from the presence of the pointlike contribution. Therefore, unlike with the proton case, an important constraint on the parton densities is missing. This constitutes another difference of the photon pdfs with those of the proton.

The general solution of the evolution equations (2.18) has an inhomogenous and a homogenous part:

$$\vec{q}^{\,\gamma} = \begin{pmatrix}\Sigma^\gamma\\ g^\gamma\end{pmatrix} = \vec{q}^{\,\gamma}_{PL} + \vec{q}^{\,\gamma}_{had} \tag{2.22}$$

The homogenous/hadronic part of the solution $\vec{q}^{\,\gamma}_{had}(x_\gamma, Q^2)$ contains a perturbatively uncalculable boundary condition $\vec{q}^{\,\gamma}_{had}(x_\gamma, Q_0^2)$. On the other hand, given a Q_0^2, the inhomogenous/pointlike component is completely perturbatively calculable. The quark densities are:

$$q_i^\gamma \sim \frac{1}{\alpha_s(Q^2)} \tag{2.23}$$

This anomalous $1/\alpha_s(Q^2)$ behaviour of the photon pdfs arises because of the inhomogenous terms coming from the direct coupling $\gamma \to q\bar{q}$.

Because of the perturbatively uncalculable part of the quark (gluon) densities, different parametrisations are used for describing the parton distribution functions in the photon.



Figure 2.6 - LO and NLO parametrizations of the real photon pdfs [Vo97]. In the above two plots are given the (up) quark densities. The two plots below show parametrizations of the gluon densities

2.3.2 Parametrizations of the parton distribution functions for real photons.

In order to calculate the photon pdfs at all scales Q^2, one needs to choose a reference scale Q_0^2 and use some Ansatz for parametrizing the pdfs at this scale. Usually a vector meson dominance (VMD) Ansatz is used for $q_i^{\gamma}(x_{\gamma}, Q_0^2)$. Then fits to the F_2^{γ} data are used to fix free parameters. Since F_2^{γ} is a sum of the quark/antiquark distributions weighted with the squared charges (see 2.15), the quark/antiquark distributions can be well determined using this method. Fixing the gluon distributions is not so easy and dependent on the Ansatz used for the initial pdfs. In lack of a momentum sum rule as for the proton, the gluon distributions cannot be directly related to the quark ones.

We know that for $Q_0^2 \geq 1$ GeV^2, a pure VMD input is insufficient for fitting the data at higher scales. Usually two methods are used to solve this insufficiency:

1. Pure VMD valence like input distributions are used and a very low scale is chosen $Q_0^2 < 1$ GeV^2 for starting the evolution (for the VMD, the valence quark distributions of the pion are used). At such a low scale the input distributions should require only a VMD component and the anomalous component can be generated dynamically from the evolution equations. The problem with this method is that we do not know whether pQCD still works at such small scales.

2. A VMD + anomalous input is used selecting a starting scale $Q_0^2 \geq 1$ GeV^2. The quark densities are fitted to the F_2^{γ}. A guess must be made about the gluon densities.

Different parametrizations in leading (LO) and next-to-leading order (NLO) of the quark and gluon distributions are shown in figure 2.6. More about parametrizations can be found in: SaS1D/SaS2D - [Sj95], GRV - [Gl92], GS - [Go92], AFG (NLO) - [Au94]. The (simplified) main characteristics of the LO parametrizations are summarized in the following table.

LO parametrization	Q_0^2 (GeV^2)	input	factorization scheme
GRV	0.25	purely VMD	DIS_{γ}
SaS1D	0.36	purely VMD	DIS_{γ}
GS	3.0	VMD + perturbative	$\overline{MS}$
SaS2D	4.0	VMD + perturbative	DIS_{γ}

The "D" of SaS1D and SaS2D parametrizations refers to the factorization scheme used - DIS_{γ}.

2.3.3 Parametrizations of the parton distribution functions for virtual photons.

Also for virtual photons different ansätze are used for parametrizing the pdfs. The hadronic (non-perturbative) VMD component is generally expected to diminish as



Figure 2.7 - Different parametrizations of the virtual photon pfd's: quarks (left plot) and gluons (right plot) at $Q^2 = 10$ GeV^2 for two different values of P^2: $P^2 = 0.2$ GeV^2 and $P^2 = 1.$ GeV^2

the virtuality (P^2) of the photon increases. Here we will explain the Ansatz used by Schuler and Sjöstrand for the pdfs of virtual photons.

For the real photon, Schuler and Sjöstrand see the pdfs as decomposed in a VMD and an anomalous component:

$$f_i^{\gamma}(x, Q^2) = \sum_{V}^{\rho^0,\omega,\phi,J/\psi} \frac{4\pi\alpha}{f_V^2} \cdot f_i^{\gamma,V}(x, Q^2, Q_0^2) \quad + \int_{Q_0^2}^{Q^2} \frac{dk^2}{k^2} \frac{\alpha}{2\pi} \sum_q 2e_q \cdot f_i^{\gamma,q\bar{q}}(x, Q^2, k^2) \tag{2.24}$$

The VMD distributions and the anomalous distributions are parametrized separately and added to give the full result. The transition region from real to virtual photons is described introducing a dipole dampening factor for each component separately:

VMD	$1/(1 + P^2/m_V^2)^2$
anomalous	$1/(1 + P^2/k^2)^2$

Additionally the lower input scale for the VMD states is shifted from Q_0^2 to $P_0^2 \approx max(P^2, Q_0^2)$.

The resulting quark and gluon densities for the SaS1D, SaS2D (and GRS which we

did not discuss here) are shown in figure 2.7 for two different values of the virtuality P^2.

A lot of measurements performed in e^+e^- experiments have helped to constrain the pdfs of the real photons by using fits to the data. Contrary to this, few experimental data exist from e^+e^- experiments to constrain the pdfs of the virtual photons. These comes mainly from the fact that double-tagged events have a very poor statistics. As we shall see in the next sections, HERA is in a better position of studying the structure of virtual photons.

2.4 Photon Structure at HERA

At HERA, protons collide with electrons (positrons). The high flux of almost on-shell photons which accompany the electron (positron) beam provides a unique opportunity to study the nature of the photon and its interactions.

2.4.1 Dijet Production in γP Interactions

The photon may interact directly with a parton from the proton or it may first fluctuate into a hadronic state. In direct interactions the photon takes part as a whole in the interaction, therefore the fraction of photon's momentum entering the hard scattering is one ($x_\gamma = 1$). In the case of as resolved interactions the photon acts as a source of partons which scatter off the partons from the proton. The quantity x_γ is now less than unity since only a fraction of the photon's momentum enters into the hard interaction. An example of two leading order pQCD diagrams of these interactions are shown in figure 2.8.



Figure 2.8 - Two example leading order diagrams for direct (left) and resolved (right) photoproduction at HERA. Note the presence of the photon remnant in the case of resolved interactions.

In both cases the final state consists of two (or more) hadron jets, the proton remnant and the scattered electron. In the case of resolved interactions, in addition to the above, the final state includes also the photon remnant produced by the spectator partons in the photon which do not contribute to the hard interaction. The photon remnant is moving in the direction of the incoming photon. The LO graphs for the direct and resolved interactions at HERA are shown in figure 2.9.

The two hard (i.e. large transverse energy) jets in the final state are related to the underlying parton dynamics. They can be used to measure the structure of the proton and the photon. Experimentally x_γ cannot be measured directly. In order to separate the resolved and direct interactions, the following observable is defined:

$$x_\gamma^{OBS} = \sum_{\text{jets}} \frac{E_T^{jet} \cdot e^{-\eta^{jet}}}{2yE_e} \tag{2.25}$$

E_T^{jet} is the transverse energy of the jet and η^{jet} is the pseudorapidity defined as:

$$\eta^{jet} = -\ln\left[\tan\left(\frac{\theta^{jet}}{2}\right)\right] \tag{2.26}$$

The sum in 2.25 runs over the two highest transverse energy jets

In the formulas (2.25,2.26) the positive z-direction is in the direction of the incoming protons. θ^{jet} is the polar angle of the jet with respect to the positive z-direction (for a more detailled definition of this angle see section on jets). Bearing in mind the correlation of the jets with the underlying partons, at LO the quantity x_γ^{OBS} is the fraction of the photon's momentum entering the hard interaction.

In figure 2.10 the variable x_γ^{OBS} is plotted. The solid line is the Herwig5.9 MC and the shaded area is the direct-only part from the simulation. One can see that a rather clean separation between the resolved and direct classes of events can be done by cutting at $x_\gamma^{OBS} = 0.75$.

At HERA there is only one photon as opposed to two photons in the case of e^+e^- experiments. By Q^2 here one denotes the virtuality of the (target) photon[3].

For high enough virtualities of the photon, the photon structure should be completely suppressed and the photon will act as a pointlike particle which probes the proton. But at smaller virtualities the photon itself will show a structure.

At HERA the photon structure is probed by a parton from the proton, i.e. by a strongly interacting object. Both quarks and gluons contribute to the interactions in ep collisions (in e^+e^- gluons do not contribute at leading order). Therefore, at HERA we are sensitive to the gluon content of the photon. Due to the absence of a simple momentum sum rule, the gluon densities cannot be extracted from the F_2^γ measurements. In jet photoproduction there is a direct sensitivity to the gluon density in the photon. This sensitivity is concentrated in the low x_γ (x_γ^{OBS}) region.

HERA is in a unique position of studying the virtual photons and the photon structure as the virtuality of the photon increases. As we have already described, such

[3]As a reminder: in e^+e^- experiments the virtuality of the target photon was denoted by P^2.



Figure 2.9 - LO graphs for direct and resolved photon interactions. The two graphs of the first (top) row: QCD Compton scattering and Boson-Gluon Fusion are the two (LO) direct photon interactions. The other graphs represent the 6 possible (LO) resolved photon interactions.



Figure 2.10 - The x_γ^{OBS} variable for dijet photoproduction events at HERA. The solid line is the Herwig5.9 simulation and the shaded area is the direct-only part from the simulation. The direct part is concentrated at higher x_γ^{OBS} and the resolved part at lower x_γ^{OBS}. One can see that a rather clean separation of the resolved and direct events can be done by a cut on $x_\gamma^{OBS} = 0.75$.

measurements in e^+e^- are difficult and suffer from poor statistics. Moreover, the photon can be probed at higher scales than in the e^+e^- experiments due to the high center-of-mass energy at HERA.

General descriptions on the photon structure at HERA can be found in [Ab98], [Ca95], and [Bu95].

The ZEUS and H1 collaborations have studied the structure of real and virtual photons. More on these studies can be found in [H1r], [H1v], [ZEr], [ZEv].

2.4.2 The two Scale Problem

In the case of $\gamma^{(*)}P$ interactions we must take into account two hard scales: the virtuality of the photon (Q^2) and the scale of the hard scattering leading to the production of jets. This second hard scale can be evaluated by the transverse energy of the jets (E_T^2).



Figure 2.11 - Photon structure seen in the phase space of the two scales at HERA. In the picture also the line is drawn where $E_T^2 =$ photon virtuality. In reality the transitions between the different regions are smooth and there are no sharp limits between them.

If the virtuality of the photon is smaller than a certain scale k_0^2, then the photon can be seen as a hadron and (as we have seen above) a VMD description of it is appropriate. The $\gamma^{(*)}P$ interaction can be seen as an VP interaction of the vector meson V with the proton P. At higher Q^2, if the photon virtuality Q^2 is smaller than the scale E_T^2, the photon will still develop a partonic structure [Fi93] coming from the anomalous photon structure. If the virtuality of the photon is larger than k_0^2 and larger than the other hard scale (E_T^2), than the photon cannot be resolved and will act as a pointlike particle. In reality there is no clear cut between the VMD-direct and direct-anomalous and the transitions between the different classes of interactions are smooth. A schematic view in the phase space of the two hard scales is given in figure 2.11. The anomalous and the VMD classes together form the class of resolved photon interactions.

2.4.3 Dijet Cross Sections - The Flux of Real and Virtual Photons

The ep interaction can be seen as two subsequent processes: emission of a photon by the electron and photon-proton interaction. If one denotes by μ the hard scale of the interaction, then according to the factorization theorem, the ep dijet cross section at HERA can be written as:

$$d\sigma\,(ep \to 2J + e + X) = \left[f_{i/\gamma}(x_\gamma, \mu) \otimes M_{i,j}(\mu, \hat{s}) \otimes f_{j/p}(x_p, \mu) \right] \cdot f_{\gamma/e}(y, Q^2) \quad (2.27)$$

$\hat{s}$ is the c.m. energy of the particles i, j. $f_{i/\gamma}$ is the density of the parton (quark, gluon, photon) i in the photon. $f_{j/p}(x_p, \mu)$ is the density of the parton j in the proton. $M_{i,j}$ is the cross section for the $2 \to 2$ parton-parton hard scattering. $f_{\gamma/e}(y, Q^2)$ is the flux of photons from the electron.

The flux of the (almost) real photons from unpolarized electrons (positrons) is given by the *Weizsäcker Williams Approximation* (or EPA - *Equivalent Photon Approximation*) [We34]. According to this approximation, the probability that an electron radiates a photon with given virtuality Q^2 and fractional energy z (in the rest frame of the proton this is the same as the inelasticity y) is given as:

$$f_{\gamma/e}(z, Q^2) = \frac{\alpha}{2\pi Q^2} \left[\frac{1 + (1-z)^2}{z} - 2\frac{1-z}{z} \cdot \frac{Q^2_{min}}{Q^2} \right] \quad (2.28)$$

where Q^2_{min} is the lowest virtuality that can be obtained. This lowest virtuality is determined from the mass of the electron m_e:

$$Q^2_{min} = m_e^2 \frac{z^2}{1-z} \quad (2.29)$$

For photoproduction events (quasi-real photons) the photon spectrum can be derived by integrating the EPA from Q^2_{min} to Q^2_{max}, where Q^2_{max} is the maximal value of the photon virtuality achievable:

$$f_{\gamma/e}(z) = \frac{\alpha}{2\pi} \left[\frac{1 + (1-z)^2}{z} \cdot \ln \frac{Q^2_{max}}{Q^2_{min}} - 2\frac{1-z}{z} \cdot \left(1 - \frac{Q^2_{min}}{Q^2_{max}} \right) \right] \quad (2.30)$$

The above formula gives a typical soft Bremsstrahlung spectrum.

The flux of the transversely polarized virtual photons emitted from the electron is [Ha84]:

$$f^T_{\gamma^*/e}(y, Q^2) = \frac{\alpha}{2\pi Q^2} \left[\frac{1 + (1-y)^2}{y} - 2\frac{1-y}{y} \cdot \frac{Q^2_{min}}{Q^2} \right] \quad (2.31)$$

and for longitudinal virtual photons:

$$f^L_{\gamma^*/e}(y, Q^2) = \frac{\alpha}{2\pi Q^2} \left[2\frac{1-y}{y} \right] \quad (2.32)$$

The above flux of longitudinally polarized photons is expected to be small.

2.5 Electromagnetic Photon Structure

The electromagnetic (or QED) photon structure is related to lepton pair production in the $\gamma^*\gamma$ scattering. The investigation of the QED photon structure is of interest because of two reasons [Ni00]:

1. it is a test of QED of the order $O(\alpha^4)$.

2. these measurements done in a clean experimental situation are used to refine the experimentalist's tools and investigate the possibilities of extracting similar information in the case of a much more complex hadronic final state.

In contrast to the hadronic structure functions discussed above, the QED structure functions $F_2^{\gamma(QED)}$, $F_1^{\gamma(QED)}$, $F_L^{\gamma(QED)}$ can be calculated in QED and all the particles in the final state can, in principle, be observed. The structure function $F_L^{\gamma(QED)}$ is, similarly to F_L^γ, negligible because of the factor y^2 in its contribution to the cross-section. Only a muonic final state allows for the extraction of the structure functions from the data. In the case of muons, the invariant mass of the $\mu^+\mu^-$ pair can be determined accurately, and hence x_{BJ}.

A summary of the $F_2^{\gamma(QED)}$ measurements can be seen in figure 2.12. These measurements are another proof of the correctness of QED.

A more detailed description of the electromagnetic structure functions can be found in [Ni00] and [Kr00].



Figure 2.12 - Summary of $F_2^{\gamma(QED)}$ measurements. The errors plotted are only statistical. The structure function $F_2^{\gamma(QED)}$ is plotted versus x_γ^{OBS} in different bins of the photon virtuality (given in brackets in each region). [Ni00]

Chapter 3

HERA and ZEUS

3.1 The Hadron Electron Ring Accelerator

Figure 3.1 - Aerial view of DESY and surroundings together with the two largest accelerators HERA and PETRA.

The Hadron Electron Ring Anlage (HERA) is a lepton proton collider located at DESY in Hamburg, Germany. It offers unique opportunities to explore the ep interactions as the only ep collider in the world. HERA was built during the late 80's and