# A search for QCD-instantons in deep-inelastic $ep$ scattering with the ZEUS detector at HERA

Dissertation
zur Erlangung des Doktorgrades
des Fachbereichs Physik
der Universität Hamburg

vorgelegt von

Sonja Hillert

aus Hamburg

Hamburg
2002

| | |
|---|---|
| Gutachter der Dissertation: | Prof. Dr. E. Lohrmann<br>Prof. Dr. R. Klanner |
| Gutachter der Disputation: | Prof. Dr. E. Lohrmann<br>Prof. Dr. V. Blobel |
| Datum der Disputation: | 20. Dezember 2002 |
| Vorsitzender des Prüfungsausschusses: | Prof. Dr. F.-W. Büßer |
| Vorsitzender des Promotionsausschusses: | Prof. Dr. G. Huber |
| Dekan des Fachbereichs Physik: | Prof. Dr. F.-W. Büßer |

## Abstract

A search for QCD-instanton induced events in deep-inelastic *ep* scattering (DIS) has been performed with the ZEUS detector at the HERA accelerator, using a neutral current sample of 1996/1997 data corresponding to an integrated luminosity of $38.6\,\mathrm{pb}^{-1}$. A kinematic range defined by cuts on the photon virtuality, $Q^2 > 120\,\mathrm{GeV}^2$, and on the Bjorken scaling variable, $x > 10^{-3}$, has been investigated. QCD-instanton induced events were modelled by the Monte Carlo generator QCDINS, predicting an instanton contribution to the inclusive sample of $\approx 0.5\%$. The background from standard DIS was described by a combination of Djangoh (Ariadne) with diffractive Rapgap and by Herwig, respectively. Different methods to enhance the fraction of instanton induced events were tried, Fisher's algorithm proving to be the best one. In an instanton-enhanced subsample, sphericity has been fitted by a combination of signal and background Monte Carlo, yielding an upper maximum likelihood $2\sigma$ limit on the fraction of instanton-induced events in the enhanced sample of 12% (8.1%) when Herwig (Djangoh with Rapgap) is assumed to describe the background correctly, compared to an expected fraction of 8.8%. However, there are indications that the simulation of standard DIS events is not reliable. Therefore, a most conservative $2\sigma$ limit of 206 instanton-induced events in the enhanced sample has been set, assuming the background to be zero, where 62 events are predicted by QCDINS.



## Kurzfassung

Mit dem ZEUS-Detektor am HERA-Beschleuniger wurde eine Suche nach QCD-Instanton induzierten Ereignissen in tiefunelastischer *ep*-Streuung (DIS) durchgeführt. Der dabei verwendete Datensatz von Neutralstrom-Ereignissen aus den Jahren 1996/1997 entsprach einer integrierten Luminosität von $38.6\,\mathrm{pb}^{-1}$. Untersucht wurde der durch Schnitte auf die Photonvirtualität, $Q^2 > 120\,\mathrm{GeV}^2$, und auf die Bjorken-Skalenvariable, $x > 10^{-3}$, definierte kinematische Bereich. Instanton-induzierte Ereignisse wurden durch den Monte Carlo Generator QCDINS beschrieben, der einen Instanton-Beitrag von ca. 0.5% zum inklusiven Datensatz vorhersagt. Der Standard-DIS Hintergrund wurde zum einen durch eine Kombination von Djangoh(Ariadne) mit diffraktivem Rapgap, zum anderen durch Herwig beschrieben. Von verschiedenen Verfahren zur Anreicherung Instanton-induzierter Ereignisse erwies sich der Fisher-Algorithmus als am besten geeignet. In einem angereicherten Datensatz wurde eine Kombination von Signal- und Hintergrund-Monte Carlo an die Sphärizitätsverteilung angepasst. Die Anpassung ergab eine obere Maximum Likelihood $2\sigma$-Grenze auf den Anteil Instanton-induzierter Ereignisse im angereicherten Datensatz von 12% (8.1%), wenn angenommen wird, daß Herwig (Djangoh mit Rapgap) die Daten korrekt beschreibt, verglichen mit einem vorhergesagten Anteil von 8.8%. Es gibt allerdings Hinweise, daß die Simulation von DIS-Ereignissen unzuverlässig ist. Daher wurde unter der Annahme, daß der DIS-Hintergrund keinen Beitrag liefert, eine konservative $2\sigma$-Grenze von 206 Instanton-induzierten Ereignissen im angereicherten Datensatz gesetzt, für den QCDINS 62 Ereignisse vorhersagt.

# Contents

# List of Figures

# Introduction

In the Standard Model, both the strong and the electroweak interaction are described by non-abelian gauge theories. In such theories, the ground state has a rich topological structure. Its simplest building blocks are non-perturbative fluctuations of the gauge fields [Bel75], called instantons ['t Hooft]. They can be interpreted as tunnelling processes of the gauge fields between topologically distinct types of vacuum states. The name instanton refers to the fact, that these fluctuations are localised in space and Euclidean time.

In quantum chromodynamics (QCD), the theory of strong interactions, a number of theoretical problems can be addressed by taking instanton effects into account, for a review see e.g. [Schä98]. For example, instanton theory predicts the non-conservation of chirality and provides an explanation for the fact that the $\eta'$ mass is much larger than that of the $\eta$ state.

Although instantons are theoretically required by the Standard Model, they have not yet been observed experimentally. While in electroweak interactions instantons might play a role only at cm energies of $\gg 10\,\mathrm{TeV}$ [Rin91, Gib95, Rin94, Rin02], in QCD instanton effects are expected to become sizable at much lower energies, because the strong coupling $\alpha_s$ is much larger than the equivalent parameter, $\alpha$, in electroweak theory.

QCD instantons are predicted to have short distance implications [Bal93, Rin94, Moch97, Rin98]. In particular, they can induce characteristic events in neutral current (NC) deep-inelastic *ep* scattering (DIS) at HERA. The cross section of these events has been calculated by A. Ringwald, F. Schrempp and collaborators using instanton-perturbation theory [Moch97, Rin98]. Besides, the phenomenology of instanton-induced events at HERA has been studied [Rin94, Car99, Rin00]: such events are characterised by a high multiplicity final state of large transverse momentum, large sphericity, flavour democracy and maximal chirality violation.
The Monte Carlo (MC) generator QCDINS [Rin00] by A. Ringwald and F. Schrempp allows dedicated experimental searches for instanton-induced events to be performed. Results of such a search have recently been reported by the H1 collaboration [Ad02]. Upper limits for the cross section of 60 pb and 100 pb were determined, depending on the kinematic region considered.

In this thesis, a ZEUS analysis using a sample of 1996/1997 neutral current data corresponding to an integrated luminosity of $38\,\mathrm{pb}^{-1}$ is presented. As instanton-induced events are predicted to contribute only $\approx 0.5\%$ to the cross section of the inclusive sample used in this search, it is essential to find cuts that efficiently reject background events, thus enhancing the instanton contribution in the subsample. For that purpose, variables characterising the hadronic final state are studied. In one attempt, combinations of one- and two-dimensional cuts are used for restricting the sample. In another approach, first an optimal linear combination of the input variables is determined using the Fisher algorithm [Fish36, BlLo98]. The enhancement is then achieved by a cut on the resulting Fisher discriminant.

Upper limits on the fraction of instanton-induced events within instanton-enhanced samples are set using two different approaches: the fit approach extracts the information on the instanton contribution from the shape of characteristic variables. In the cut approach a limit is determined from the number of events passing a set of hard cuts.

# Chapter 1

# Theoretical Background

## 1.1 Electron-proton scattering




Figure 1.1: *A generic diagram of ep scattering. Four-momenta of the interacting particles are given in parentheses.*

In electron-proton scattering, a point-like particle, the electron[1], interacts via the electromagnetic or weak force with the proton, which has a complex substructure. The interaction can be described by the exchange of a photon, a $Z^0$ boson or a $W^{\pm}$, transferring a four-momentum $q$ from the lepton to the proton as shown in Fig. 1.1.
Neutral current (NC) processes are mediated by a photon or $Z^0$, while $W^{\pm}$ exchange gives rise to charged current (CC) events. The contributions of NC and CC events to the total cross section depend on the virtuality $Q^2 = -q^2$ of the exchanged boson. While the NC cross section dominates at medium $Q^2$ due to the dominance of the electromagnetic over the weak contribution, at virtualities $Q^2 \approx 10^4\,\mathrm{GeV}^2$ the NC and CC cross sections are of similar size, see Fig. 1.2. This is evidence that the electromagnetic and the weak contributions can be described by a unified "electroweak" field theory, in which the couplings of the two contributions obtain similar values in that high $Q^2$

[1]In what follows, both positrons and electrons are referred to as electrons.




Figure 1.2: *Comparison of Neutral Current and Charged Current cross sections as a function of $Q^2$ measured in deep-inelastic ep scattering at HERA. From [Zha02].*

range.

For the analysis presented in this thesis, a NC deep-inelastic scattering (DIS) sample at $Q^2$ values, at which $Z^0$ exchange can be neglected, has been used. In that regime, the photon acts as a probe to study the substructure of the proton and the nature of the strong force giving rise to it. All processes observed in DIS, except for instanton-induced events, will be referred to as normal DIS (nDIS) events in what follows.

The Born cross section can be expressed in terms of the proton structure functions $F_i^{\rm NC}$, $i = 1, 2, 3$, and the kinematic variables $x$, $y$ as

$$\left(\frac{d^2\sigma^{\rm NC}}{dx\; dQ^2}\right)_{\rm Born} = \frac{4\pi\alpha^2}{x\,Q^4}\left\{y^2 x F_1^{\rm NC}(x,Q^2) + (1-y)F_2^{\rm NC}(x,Q^2) \mp \left(y - \frac{y^2}{2}\right) x F_3^{\rm NC}(x,Q^2)\right\} .$$

Two structure functions, $F_1$ and $F_2$ are needed to describe the interaction of the photon with the proton. The combination

$$F_L(x,Q^2) = \left(1 + \frac{4M^2x}{Q^2}\right) F_2(x,Q^2) - 2xF_1(x,Q^2) \approx F_2 - 2xF_1 \quad ,$$

is the longitudinal structure function, where $M$ is the proton mass. The function $F_1$ ($F_L$) corresponds to the absorption of transversely (longitudinally) polarised photons. The function $xF_3^{\mathrm{NC}}$ accounts for parity violation due to $Z^0$ exchange, hence contributing only at large $Q^2$.
The kinematic variables $Q^2$, $x$ and $y$ are defined as

$$Q^2 = -q^2 = -(k-k')^2 \quad , \qquad x = \frac{Q^2}{2P \cdot q} \quad , \qquad y = \frac{q \cdot P}{k \cdot P} \quad \text{(cf. Fig. 1.1)} \quad .$$

Because of the relation $Q^2 = x \cdot y \cdot s$, with $s$ being the $ep$ cm energy squared, the kinematics of the process is fully determined by any choice of two of the three variables $x$, $y$, $Q^2$.
The variables $x$ and $y$ are interpreted in the quark-parton model, which states that an $ep$ scattering process can be approximated by the scattering of the electron off a quark inside the proton. In that model, the Bjorken variable $x$ is the fraction of the proton momentum carried by the struck parton, while $y$ represents the fraction of the positron energy transferred to the proton in the rest frame of the initial-state proton. The structure functions of the proton at large values of $Q^2$ have been precisely measured at HERA [StrFct].

## 1.2 Quantum Chromodynamics

The strong force is described by quantum chromodynamics (QCD), a renormalisable, non-abelian gauge field theory, see e.g. [Ell96] and the references therein. It is based on an $SU(3)$ symmetry between three types of colour charge carried by the quarks. However, because QCD is non-abelian, its gauge bosons, the gluons, also carry colour and can therefore interact with each other. The strength of the interaction is determined by the coupling $\alpha_s$, which is not constant in a renormalisable gauge theory, but depends on the energy scale, usually chosen as $Q$.
At low energies, or equivalently, large distances, the coupling is large. Therefore, quarks and gluons cannot be observed as free particles, but are confined within colour-neutral composed objects, such as the proton. At those scales, interactions can only be described non-perturbatively. However, at large scales (short distances), $\alpha_s$ is small enough for the application of perturbation theory.
Physical observables $R(Q)$ can then be written in terms of $\alpha_s$ as

$$R(Q) = R_0(Q) + \alpha_s R_1(Q) + \alpha_s^2 R_2(Q) + \ldots \quad .$$

They should not depend on the renormalisation scale $\mu_R$, at which ultraviolet divergences are subtracted, as $\mu_R$ can be chosen arbitrarily, so

$$\mu_R^2 \frac{d}{d\mu_R^2} R\left(\frac{Q^2}{\mu_R^2},\ \alpha_s\right) = 0 \quad .$$

If the renormalisation scale is chosen such that both $\alpha_s(Q)$ and $\alpha_s(\mu_R)$ are small enough, the energy scale dependence of $\alpha_s$ can be determined within the framework of perturbation theory. To lowest order it is given by [Ell96]

$$\alpha_s(Q^2) = \frac{\alpha_s\left(\mu_R^2\right)}{1+\alpha_s\left(\mu_R^2\right)bt} \quad , \; t = \ln\frac{Q^2}{\mu_R^2} \quad , \; b = \frac{1}{4\pi}\left(11 - \frac{2}{3}n_f\right) \quad ,$$

where $n_f$ is the number of quark flavours with masses below $\mu_R$.
The value of $\alpha_s$ at a fixed scale, usually given by the mass of the $Z$ boson, $M_Z$, can be determined in various ways, for an overview see e.g. [Beth00]. At HERA, $\alpha_s$ has been determined from jet physics [AlS], in particular from inclusive jet cross sections and dijet cross sections, as well as from measurements of the jet shape and the subjet multiplicity. Results are found to be in agreement with measurements from $p\bar{p}$ [Af02] and $e^+e^-$ interactions, see [Beth00], and with the current world average [PDG00] of

$$\alpha_s(M_Z^2) = 0.1181 \pm 0.0020 \quad .$$

Both perturbative and non-perturbative effects have to be accounted for in the description of deep-inelastic $ep$-scattering. The factorisation theorem [Col85] states that the scattering process factorises with respect to these contributions, such that the hard scattering process, calculable in perturbative QCD, can be treated independently from non-perturbative effects. These are absorbed into universal parton distribution functions of the proton that have to be determined experimentally.
Three types of hard scattering processes contribute to the neutral current cross sec-




(a) (b) (c)

Figure 1.3: *Processes contributing to the NC cross section up to $\mathcal{O}(\alpha_s)$: a) Quark-Parton-Model, b) QCD-Compton, c) Boson-Gluon-Fusion.*

tion up to $\mathcal{O}(\alpha_s)$: the quark-parton model process (QPM), the QCD Compton process (QCD-C) and the boson-gluon fusion (BGF), shown in Fig.1.3.
The structure functions $F_i(x, Q^2)$ of the proton, introduced in section 1.1, are related to the partonic structure functions $\hat{F}_i^a$, which describe the hard scattering process between the virtual photon and parton $a$ ($a$ being a gluon, quark or antiquark) via the convolution

$$F_i(x, Q^2) = \sum_a f_a(x, \mu_F) \otimes \hat{F}_i^a\left(x, \frac{Q}{\mu_F}, \alpha_s(\mu_R)\right) \quad ,$$

with the parton density functions $f_a(x, \mu_F)$. These expressions depend on the factorisation scale $\mu_F$, which is commonly set equal to the renormalisation scale: $\mu_F = \mu_R = Q$. In addition to the factorisation scale dependence, $f_a$ and $\hat{F}_i^a$ depend on the factorisation scheme, which determines, how much of the finite contribution to $F_i$ is absorbed into the parton densities. Two schemes commonly used are the minimal subtraction ($\overline{MS}$) scheme and the DIS scheme.
The parton densities $f_a$ are themselves scale dependent. However it is sufficient to experimentally determine them at an initial scale, as their evolution to different scales is predicted by QCD. That evolution is described by a set of $(2n_f - 1)$ equations referred to as the DGLAP equations, derived by Dokshitzer, Gribov, Lipatov, Altarelli and Parisi [DGLAP]. Using $t = \mu^2$, it can be written in the form

$$t\frac{\partial}{\partial t}\begin{pmatrix} q_i(x,t) \\ g(x,t) \end{pmatrix} = \frac{\alpha_s(t)}{2\pi}\sum_{q_i,\bar{q}_j}\int_x^1 \frac{d\xi}{\xi}\times\begin{pmatrix} P_{q_iq_j}\left(\frac{x}{\xi},\alpha_s(t)\right) & P_{q_ig}\left(\frac{x}{\xi},\alpha_s(t)\right) \\ P_{gq_j}\left(\frac{x}{\xi},\alpha_s(t)\right) & P_{gg}\left(\frac{x}{\xi},\alpha_s(t)\right) \end{pmatrix}\begin{pmatrix} q_i(\xi,t) \\ g(\xi,t) \end{pmatrix} ,$$

where $q_i(x, t)$ and $g(x, t)$ are the (anti-)quark and gluon densities, respectively. The functions denoted by a capital $P$ with quark/gluon indices are the splitting functions, each of which is calculable as a power series in $\alpha_s$,

$$\begin{aligned} P_{q_iq_j}(z,\alpha_s) &= \delta_{ij}P^{(0)}_{qq}(z) + \tfrac{\alpha_s}{2\pi}P^{(1)}_{qq}(z) + \ldots \\ P_{q_ig}(z,\alpha_s) &= \delta_{ij}P^{(0)}_{qg}(z) + \tfrac{\alpha_s}{2\pi}P^{(1)}_{qg}(z) + \ldots \\ P_{gq_j}(z,\alpha_s) &= \delta_{ij}P^{(0)}_{gq}(z) + \tfrac{\alpha_s}{2\pi}P^{(1)}_{gq}(z) + \ldots \\ P_{gg}(z,\alpha_s) &= \delta_{ij}P^{(0)}_{gg}(z) + \tfrac{\alpha_s}{2\pi}P^{(1)}_{gg}(z) + \ldots \quad . \end{aligned}$$

They describe the probability of parton emission by quarks and gluons, depending on the types of partons involved, and on the distribution of the initial momentum to the resulting partons. As a consequence of charge conjugation invariance and $SU(n_f)$ flavour symmetry, the splitting functions are the same for quarks and antiquarks and independent of the quark flavour.

## 1.3 Instantons

Instantons are the simplest building blocks of the rich topological structure, which forms the ground state of QCD. Being a non-abelian gauge theory, QCD gives rise to topologically distinct vacuum states, which are separated from each other by finite energy barriers [Rin94]. However, the gauge fields may tunnel between the different states. These rapid tunnelling events, localised in space and Euclidean time, are described by instantons.
The presence of an instanton influences the hard subprocess, which forms the core of deep-inelastic *ep*-scattering events. As developed theoretically and phenomenologically by A. Ringwald and F. Schrempp [Rin94, Moch97, Rin98, Rin99, Rin00, Rin01], it results in a characteristic final state, that allows instanton-induced events to be distinguished from events stemming from normal DIS ("nDIS") processes. The Monte Carlo

generator QCDINS [Rin00], describing instanton-induced events in the HERA regime, predicts a sizable rate of these events. To the experimentalist, HERA therefore offers a unique chance to discover instantons, which would be a confirmation of an essential non-perturbative Standard Model prediction.

### 1.3.1 A quantum mechanical example




Figure 1.4: *The double well potential for (a) Minkowski time and (b) Euclidean time $\tau$. In (a) the dots at $x = \pm x_0$ represent the classical solutions. In (b), the region $-x_0 < x < x_0$ becomes accessible. Figure from [Schr99].*

Some basic properties of instantons can be studied in the context of classical quantum mechanics. Consider the double-well potential with minima at $\pm x_0$, $V(x) = g(x^2 - x_0^2)^2$, illustrated in Fig. 1.4(a). Classically, the ground state has the two degenerate solutions $x = \pm x_0$. Quantum mechanically, tunnelling between these states can occur. Although being a quantum mechanical process, this can be described in a classical way. The classical equation of motion is given by

$$E = \frac{1}{2}\left(\frac{dx}{dt}\right)^2 + V(x) \quad .$$

By replacing time $t$ by Euclidean time $\tau$: $t \to -i\tau$, one obtains

$$-E = \frac{1}{2}\left(\frac{dx}{d\tau}\right)^2 - V(x)$$

instead. So the relative sign between the potential and the time derivative of $x$ changes and the interval $-x_0 < x < x_0$ becomes accessible, cf. Fig. 1.4(b). In particular, for




Figure 1.5: *(a): instanton solution for the double well potential, (b): the solution is located at $\tau_0$. Figure adapted from [Schr99].*

$E = 0$ one finds the solution

$$x^{(\mathrm{I})}(\tau) = \pm x_0 \tanh\left(\sqrt{2g}x_0\,(\tau - \tau_0)\right) \quad ,$$

plotted in the $x - \tau-$plane in Fig. 1.5(a). This solution, the instanton (+) or anti-instanton (−) of the double-well potential, describes tunnelling between the classically degenerate ground states $\pm x_0$, hence lifting the degeneracy. Fig. 1.5(b) shows the time derivative of $x$ as a function of Euclidean time $\tau$. The distribution is sharply peaked at $\tau_0$: the instanton solution is localised in (Euclidean) space-time. Tunnelling occurs very rapidly, almost instantaneously.

The double-well potential may also be treated using the Feynman path integral method. Because that approach can be directly transferred to the mathematically more subtle case of QCD, it will be outlined in the following. The path integral method allows the probability amplitude for the transition from initial state $x(t_i) = x_i$ to final state $x(t_f) = x_f$ to be calculated as an integral over all paths $[Dx]$ between these states, weighted with their action $S(x)$:

$$\langle x_f \left|\exp\left(-iH\left(t_f - t_i\right)\right)\right| x_i\rangle = \mathcal{N}\int_{x_i}^{x_f}[Dx]\exp\left(iS(x)\right) \quad ,$$

where $H$ is the Hamiltonian and $\mathcal{N}$ a normalisation factor.
For Euclidean time this turns into

$$\langle x_f \left|\exp\left(H\left(\tau_f - \tau_i\right)\right)\right| x_i\rangle = \mathcal{N}\int_{x_i}^{x_f}[Dx]\exp\left(-S_E(x)\right) \quad ,$$

where the Euclidean action $S_E(x)$ is obtained from the one in Minkowski space as

$$S = \int dt L\left(x, \dot{x}\right) \Rightarrow iS_E \quad .$$

It depends on the Lagrangian, which for the double-well potential can be written as

$$L\left(x, \frac{dx}{dt}\right) = \frac{1}{2}\left(\frac{dx}{dt}\right)^2 - V(x) \quad .$$

That the action is finite shows that the instanton solution has a finite transition probability. For the path integral method, the classical (instanton) solutions correspond to the paths with minimal action in Euclidean space-time, which dominate the path integral. In Minkowski space-time one would instead have to integrate over various smaller contributions, which is unfeasible in practice. So using Euclidean instead of Minkowski space-time can be thought of as choosing a reference frame suited to the problem in the sense that it yields a mathematically concise description.

### 1.3.2 Instantons in QCD

QCD is a non-abelian gauge theory. The gauge field transforms as

$$\vec{A}' = U\left(\vec{x}\right)\vec{A}U^\dagger\left(\vec{x}\right) - iU\left(\vec{x}\right)\vec{\nabla}U^\dagger\left(\vec{x}\right) \quad ,$$

with a transformation $U\left(\vec{x}\right) \in SU(3)$. The vacuum is defined to be the state with minimum field energy. Other than in QED, where just one such state exists, there are infinitely many degenerate such states in QCD due to the non-abelian nature of QCD. From the transformation rule for the gauge field one can deduce that one obtains minimal field energy not only if $\vec{A} = 0$, but for all pure gauges

$$\vec{A}_{\text{vac}} = -iU\left(\vec{x}\right)\vec{\nabla}U^\dagger\left(\vec{x}\right) \quad .$$

Further, in $SU(3)$ one can distinguish between different types of gauge transformations $U\left(\vec{x}\right)$: some of these can be continuously deformed[2] into each other, others can not. All gauge transformations, which *can* be continuously deformed into each other, form a homotopy class. There are infinitely many of these classes. The different classes can be distinguished by their topological winding number

$$n = 0, \pm 1, \pm 2, \ldots \quad .$$

Not only are there different types of gauge transformations $U\left(\vec{x}\right)$, but also different types of vacua associated with the homotopy classes: The winding number of a gauge field $A$ is given by the Chern-Simons number [Rin94]

$$N_{\text{CS}} = \frac{g^2}{16\pi^2}\int d^3x \epsilon_{\text{ijk}}\left(A_i^a \partial_j A_k^a - \frac{g}{3}\epsilon_{\text{abc}}A_i^a A_j^b A_k^c\right) \quad ,$$

[2] Rotation is an example of a continuous transformation, reflexion an example of a non-continuous one.




Figure 1.6: *The different QCD vacua, indicated by different pure gauges $A_{(\mathrm{n})}$, where $n$ is the winding number, are separated by energy barriers. The degeneracy of the vacua is lifted by instantons, tunnelling solutions between "neighbouring" vacua ($\Delta N_{\mathrm{CS}} = 1$). Figure adapted from [For00].*

where $\epsilon_{\mathrm{ijk}}$ is the totally antisymmetric tensor,

$$\epsilon_{\mathrm{ijk}} = \begin{cases} +1 & \text{for ijk an even permutation of 1, 2, 3} \\ -1 & \text{for ijk an odd permutation of 1, 2, 3} \\ 0 & \text{otherwise} \end{cases} \quad .$$

Each type of vacuum comprises all pure gauges $\vec{A}_{\mathrm{vac}}$, which are obtained from the various gauge transformations $U^{(\mathrm{n})}(\vec{x})$ of one specific homotopy class $(n)$ and is separated by energy barriers from those fields $\vec{A}_{\mathrm{vac}}$, which one obtains from a transformation $U^{(\mathrm{m})}(\vec{x})$ of any different class $(m)$. Hence one arrives at the conclusion that the ground state of QCD should be infinitely degenerate. However, this is not the case. In the same way as in the simple quantum-mechanical example, the degeneracy between the states is lifted by tunnelling, i.e. instantons. In QCD, instantons describe tunnelling between neighbouring vacua, separated from each other by only one energy barrier as depicted in Fig. 1.6 and differing by only one unit of the topological winding number $N_{\mathrm{CS}}$. The instanton is therefore assigned a topological charge $Q = \Delta N_{\mathrm{CS}} = +1$. (There are also anti-instantons with $Q = -1$.)

### 1.3.3 Instanton size and distance

The one-instanton solution of the classical Euclidean field equation in QCD can be obtained analytically [Bel75]. If the Lagrange density $\mathcal{L}\left[A_\mu^{(\mathrm{I})}(x)\right]$ is calculated with




Figure 1.7: *Lagrange density of an instanton at $x_0 = 0$, from [Schr98].*

this known instanton solution, one finds a peaked behaviour

$$\mathcal{L}\left[A_\mu^{(1)}(x,\rho)\right] = \frac{12}{\pi\alpha_s}\frac{\rho^4}{\left(x^2+\rho^2\right)^4} \quad ,$$

as depicted in Fig. 1.7 . One should note that $\mathcal{L}$ only depends on the square of the coordinate $x$ in Euclidean space-time. The instanton size $\rho$ is a measure of the width of the Lagrange density distribution. From $\mathcal{L}$ one would find the Euclidean action $S_E$ to be

$$S_E = \frac{2\pi|Q|}{\alpha_s} \quad ,$$

independent of $\rho$. It is determining the tunnelling amplitude

$$T = f\left(\alpha_s\right)\exp\left(-S_E\right)\left(1 + \text{corrections}\right)$$

and hence the cross sections

$$\sigma \propto \exp\left(-\frac{4\pi}{\alpha_s}\right) \quad .$$

One sees that the tunnelling is exponentially suppressed, as one might expect from quantum mechanics. However, due to the large QCD coupling $\alpha_s$, the QCD instanton cross section is nevertheless sizable[3]. To make predictions of experimentally observable effects of QCD instantons in DIS one is not primarily interested in the exact form of the one-instanton solution, but has to take into account that Euclidean space is filled with many instantons of various sizes at some distance $R$ from each other. This is illustrated in Fig. 1.8, which shows the Lagrange density as a function of one spatial

[3]In electroweak theory, which is also non-abelian and hence also has instanton solutions, the respective cross section is much more strongly suppressed than the one for QCD, as $\alpha \ll \alpha_s$.




Figure 1.8: *Lagrange density of a slice of space time containing several instantons (size $\rho$, distance $R$). Figure from [Chu94].*

coordinate and Euclidean time obtained from a lattice calculation [Chu94].
For the kinematic range accessible in deep inelastic *ep* scattering at HERA, the predicted effects are induced by instantons of small size compared to their distance, and hence the dilute gas approximation is valid. It is then possible to calculate the cross section of the instanton-induced hard subprocess using instanton-perturbation theory. This is performed by the Monte Carlo generator QCDINS, cf. section 3.1.

### 1.3.4 Characteristics of instanton-induced events




DIS variables:
$Q^2 = -q^2 = -(e - e')^2$
$x = Q^2/(2P \cdot q)$
$W^2 = (q + P)^2 = Q^2(1 - x)/x$
$\hat{s} = (q + g)^2$
$\xi = x(1 + \hat{s}/Q^2)$

Variables of the instanton subprocess:
$Q'^2 = -q'^2$
$x' = Q'^2/(2g \cdot q')$
$s' = (q' + g)^2 = Q'^2(1 - x')/x'$

Figure 1.9: *Kinematics of instanton-induced events at HERA, from [Kuhl97]*

Fig. 1.9 shows the kinematics of instanton-induced events at HERA. The incoming lepton emits a photon, which in turn transforms into a quark-antiquark pair. One of these hadronises to form the current jet. The other one, denoted by $q'$, fuses with a gluon from the proton. The phenomenological characteristics of instanton-induced events, as worked out in [Rin94], can be summarised as follows: In the hard subprocess exactly one $q\bar{q}$ pair of each of the $n_f$ quark flavours, which are kinematically accessible, participates in the hard subprocess of each event, see Fig. 1.10. This feature is often termed "flavour democracy". The number of gluons emitted per event follows




Figure 1.10: *The hard subprocess of instanton-induced events. Figure from [Ger98].*

a Poisson distribution with average value 3. The gluons plus $2n_f - 1$ (anti-)quarks give rise to a high multiplicity final state of high transverse energy. Particles stemming from the instanton are expected to be isotropically distributed in their centre of mass frame. Therefore, identification of events possibly caused by instantons requires a good separation of the instanton from the current jet and would profit from a good reconstruction of the instanton centre of mass frame.
The most prominent characteristic of instanton-induced events from a theoretical point of view, but also the one most difficult to observe experimentally, is chirality violation. In instanton-induced events, all quarks produced in the hard subprocess are emitted with the same handedness. This means that the axial charge $Q_5$, defined as the difference between the numbers of right handed and left handed fermions, $Q_5 = \#R - \#L$, changes by the maximal possible amount of $\Delta Q_5 = 2n_f Q$ in each event.

# Chapter 2

# The ZEUS detector at the HERA collider

## 2.1 The electron proton collider HERA

The Hadron-Elektron-Ring-Anlage (HERA) [HERA81] is the only accelerator world wide, in which electrons and protons are collided. The machine consists of two storage rings mounted on top of each other in a 6.3 km tunnel $15-25$ m underground, see Fig. 2.1.

The injection of the two beams is accomplished by a series of preaccelerators, some




Figure 2.1: *(a): the HERA storage ring and (b): its preaccelerators.*

of which already existed when HERA was planned. Electrons are accelerated in a

system of two linear accelerators, reaching energies of 220 MeV in the first (LINAC 1) and 450 MeV in the second (LINAC 2), respectively. Positrons are accumulated in the positron intensity accumulator (PIA), from which they are transferred to DESY II, where they are accelerated to an energy of 7 GeV. Up to 70 bunches, already with the HERA spacing of 96 ns, are then accelerated in PETRA II to 14 GeV, and finally injected into HERA.
The proton beam is obtained by accelerating $H^-$ ions up to 50 MeV in a linear accelerator (LINAC). After removing the electrons, the protons are consecutively injected into DESY III and PETRA, where they are accelerated to 7.5 GeV and 40 GeV, respectively.
The final increase in energy in the HERA accelerator itself yields an electron beam of 27.5 GeV and a proton beam of 820 GeV (920 GeV since 1998), corresponding to a centre of mass energy of approximately 300 GeV. The magnetic field of 4.68 T, necessary to bend the high momentum proton beam in the four curved sections of the storage ring, is supplied by superconducting magnets, operating at 4.4 K. The lifetime of the proton (positron) beam is of the order of a few days ($\approx$ 8 hours).
While up to 210 bunches of each particle type can be stored in HERA, the actual




Figure 2.2: *Luminosity delivered by HERA in the years 1993 to 1997.*

| HERA parameters | Design Values | | 1996 | | 1997 | |
|---|---|---|---|---|---|---|
| | $e^-$ | $p$ | $e^+$ | $p$ | $e^+$ | $p$ |
| **Energy** [GeV] | 30 | 820 | 27.5 | 820 | 27.5 | 820 |
| $\mathcal{L}_{\text{int}}$ [pb$^{-1}$] | 35 | | 17.6 | | 36.35 | |
| **Current** [mA] | 58 | 163 | 23 | 60 | 30 | 75 |
| **Number of bunches** | 210 | | $177 + 28$ | $177 + 3$ | $176 + 20$ | $176 + 5$ |

Table 2.1: *Some design parameters of the HERA accelerator and average values found during the '96/'97 data taking period. Both the number of colliding bunches and the number of non-colliding bunches are given. From [Kl01].*

number is lower most of the time, allowing for background studies using non-colliding bunches for beam-related background and empty bunches for background due to cosmic muons. Table 2.1 gives an overview over some HERA design parameters and the actual performance during the 1996/1997 data taking period. The integrated luminosity, $\mathcal{L}_{\text{int}} = \int \mathcal{L}\text{dt}$, has steadily been increased over the years as can be seen from Fig. 2.2.

Four experiments are located at the straight sections of the HERA storage ring: the two collider experiments H1 and ZEUS, and the fixed target experiments HERMES to study the spin structure of the nucleon and HERA-B. At HERMES, longitudinally polarised electrons are inelastically scattered by a polarised gas jet target. HERA-B uses a wire-target in the halo of the proton beam. All four experiments, as well as HERA itself, underwent a major upgrade in 2000 [HERA98] in order to increase the luminosity by a factor of five and to improve the detector performance.

## 2.2 The ZEUS detector

The ZEUS detector [ZEUS93] is an almost hermetic multipurpose detector designed to identify particles produced in *ep*-scattering events and to measure their momenta and energies. Since the detector was first commissioned in 1992 it has continuously been modified to improve its performance and to extend the range of observables measured. The ZEUS coordinate system is an orthogonal, right-handed system with the $z$-axis, the 'forward direction', being defined by the direction of the proton beam. The $x$-axis points towards the centre of the HERA-ring, the $y$-axis points upwards.
As the energies of electron and proton differ strongly, the centre of mass system of the reaction is moving with respect to the laboratory frame. The asymmetric detector design, shown in Fig. 2.3, accounts for this effect. In the data taking period considered, the innermost detector component surrounding the nominal interaction point[1] was the central track detector (CTD), described in section 2.2.1. Besides the CTD, the track-

[1]The vertex detector VXD had been removed in 1995 and was replaced by a silicon microvertex detector (MVD) in 2001.




Figure 2.3: *A cross section through the ZEUS detector*

ing system comprises the FDET in the forward- and the RTD in the rear direction. While the RTD is a planar drift chamber, the forward tracking detector consists of three parts, FTD1 (the innermost) to FTD3. In the gaps between these, transition radiation detectors (TRD) are located. The tracking system is enclosed in a solenoid providing a magnetic field of 1.43 T. Momentum and charge of particles traversing the CTD can be determined from the curvature of their tracks in this magnetic field.
Energies of both charged and neutral particles are measured in the calorimeter (CAL), cf. section 2.2.2. Before particles enter the calorimeter, they pass the presampler (PRES), one layer of scintillator tiles, which allow correcting for energy losses that are caused by showering in dead material in front of the CAL. The hadron-electron-separator (HES) consists of silicon diodes, placed 2 cm below the surface of the forward and rear part of the calorimeter in order to improve the discrimination between electromagnetic and hadronic showers for particles of energies below 5 GeV.
The calorimeter is surrounded by a yoke of 7.3 cm thick iron plates, serving as return path for the magnetic flux of the solenoid. Besides, the iron-yoke is used as absorber for the backing calorimeter (BAC). In a set of proportional chambers, the BAC measures energies of showers which extend beyond the main calorimeter. Muons pass all the sub-detectors mentioned so far. They can be observed in the muon detectors FMUI, BMUI and RMUI on the inner side of the BAC and in the FMUON, BMUON and RMUON on its outer side.
Supplementary to the central detector described so far, the ZEUS experiment extends

in both directions along the beam pipe, where several components have been added to increase the angular range of measurement. Between RTD and RCAL, the small angle rear tracking detector (SRTD) with a radius of $\approx 340\,\mathrm{mm}$ has been constructed around the beam pipe. In the RCAL beam hole, an electromagnetic beam pipe calorimeter (BPC) was installed in 1995, followed in 1997 by the beam pipe tracker (BPT), located in front of the BPC towards the main detector to improve the position resolution of the BPC.

In the forward direction, the forward plug calorimeter (FPC) was installed in 1998, extending the calorimeter coverage by one unit in pseudorapidity $\eta = -\ln \tan \theta/2$. The lead-scintillator strip detectors forming the proton remnant tagger (PRT) are located at $z = 5.1\,\mathrm{m}$ in the forward direction, covering a pseudorapidity range of $4.3 < \eta < 5.8$. The leading proton spectrometer (LPS) consists of six separate silicon strip detectors, placed into the gaps between HERA beam magnets at distances between 24 m and 90 m from the interaction point. Forward neutrons can be detected in the lead-scintillator calorimeter FNC at $z = 103.6\,\mathrm{m}$. Three small electromagnetic calorimeters, the 8 m-, 35 m- and 44 m taggers, serve to measure electrons close to the beam pipe. Also the luminosity monitor is located near the forward beam pipe, see section 2.2.3.

In the rear direction, at $z = -7.3\,\mathrm{m}$, two layers of scintillator hodoscopes surrounding an iron wall (VETO) are used to reject proton-beam related background at the trigger level. The trigger system, outlined in section 2.2.4, is distributed over the entire detector volume.

### 2.2.1 The Central Track Detector (CTD)

The central track detector (CTD) [CTD] is a cylindrical wire drift chamber. Signals are collected on sense wires, located at regular intervals, while additional wires provide gain and drift fields. The CTD active volume is filled with a gas mixture consisting of 89% argon, 5% $CO_2$ and 12% ethane. It has a length of 2030 mm with an inner radius of 190 mm and an outer radius of 785 mm, respectively, covering a range of $11.6° < \theta < 168.0°$ in the polar angle $\theta$. This is equivalent to pseudorapidities $\eta$ within $2.31 > \eta > -2.26$.

Charged particles traversing the CTD ionise the gas along their path, producing $10 - 20$ ions/ cm for minimum ionising particles. With a drift velocity of about $50\,\mu\mathrm{m/\,ns}$, the freed electrons move towards the sense wires, where a stronger field gives rise to an increase in the number of electrons by a factor of $\approx 10^4$, hence yielding a measurable signal. Sense wires are read out every 9.6 ns with 8 bit precision. The timing information is used to reconstruct the hit position, which for a single hit is ambiguous, as it may be located on either side of the wire. Only the track reconstruction, using information from many wires, allows one to distinguish between real and ghost hits.

Sense wires are arranged in 9 superlayers, each consisting of 8 layers. Within each superlayer, 8 sense wires form a cell, along with the wires necessary to provide the gain and drift fields. Altogether, the CTD contains 576 cells. One octant of the CTD is